

## ***Racomitrium elongatum* FRISVOLL, a neglected Slovakian bryophyte**

**RUDOLF ŠOLTÉS**

Institute of High Mountains Biology, 059 56 Tatranská Javorina, Slovakia, e-mail:  
soltes@uniza.sk

---

Šoltés R. (2008): *Racomitrium elongatum* FRISVOLL, a neglected Slovakian bryophyte. – Thaiszia – J. Bot. 18: 59-64. – ISSN 1210-0420.

Abstract: Chorology, ecology and phytocoenology of *Racomitrium elongatum* Frisvoll is dealt with in the paper. The taxon is neglected in Slovakia and appears to be rather common in The Tatra Mts. The distribution map of the species in the Tatra Mts is presented.

Keywords: *Racomitrium elongatum*, The Tatra Mts, bryophytes, Slovakia.

---

### **Introduction**

*Racomitrium elongatum* FRISVOLL hasn't been still recorded in Slovakia. The species is a member of the *Racomitrium canescens* (HEDW.) BRID. group with a boreal distribution (DÜLL 1994). *R. elongatum* (Fig. 1) differs from *R. canescens* (Fig. 2) by having leaves strongly keeled above, straight nerve ending in apex and hair point decurrent down margin (Fig. 3). *Racomitrium canescens* leaves are either not keeled or only bluntly keeled above, the nerve is usually forked above- extending to 1/2 – 3/4 way up leaf and the hair point is rather wide, not decurrent down margin.

The distribution map of *Racomitrium elongatum* on a world scale was published by BEDNAREK-OCHYRA (1995). There are two main centres of occurrence in America – the Pacific part of the continent and Ontario. More limited distribution are found in New Foundland and Labrador.

In Europe, the moss occurs abundantly in Iceland, Faroe Islands, West and South Fennoscandia – in areas of wet, oceanic climate. Also, *Racomitrium elongatum* is frequently found in the British Islands, Portugal, North Spain, France, Netherlands, Germany, and northern Poland. It has a scattered

distribution in The Carpathians and The Alps with some isolated locations in the Balkan Peninsula, Apennine Peninsula, East and South Greenland and in Sicily. The most southerly location is in Madeira (BEDNAREK-OCHYRA, 1995).




Fig. 1. *Racomitrium elongatum*, leaf



Fig. 2. *Racomitrium canescens*, leaf




Fig. 3. *Racomitrium elongatum*, hair point decurrent down margin

In Poland, the occurrence is bicentric – Pomerania, which has an oceanic climate, and West Beskydes and The Tatras. In the Tatras, the moss reaches its highest altitude of 1700 m a. s. l. (BEDNAREK-OCHYRA 1995).

In Czech Republic, there are only poor information on the occurrence of the species. It is supposed, the habitats and the incidence of the species are similar to the species *R. canescens*. In lowland occurs rarely, more often is met in mountains (VÁŇA et al. 2008).

OCHYRA & BEDNAREK-OCHYRA (2004) placed the species with tall, conical papillae (including *Racomitrium elongatum*) to the genus *Niphotrichum*.

The 3rd species of this *Racomitrium* group is *R. ericoides*. This north-oceanic species occurs very rarely in the Tatra Mts in high-altitude level both on granite and limestone bedrocks.

## Material and methods

Collection of *Racomitrium* specimens stored in the Museum of the Tatra National Park in Tatranská Lomnica, Slovakia, was submitted to revision. The plants were collected in 1954 – 2004.

The nomenclature follows KUBINSKÁ & JANOVIČOVÁ (1998), or FRISVOLL (1983) respectively. All the specimens are stored in the Museum of the Tatra National Park in Tatranská Lomnica, Slovakia.

## Results and discussion

Revision of *Racomitrium* specimens stored in the Museum of the Tatra National Park in Tatranská Lomnica, has identified the following locations of *Racomitrium elongatum* in the Tatra Mts (Fig. 4):

1. The High Tatra Mts, Velická dolina valley, Kvetnica, granite rock, 1850 m a. s. l., leg. Šmarda, August 1954, specimen Nr.1/00205
2. The Belanian Tatra Mts, Dolina Siedmich prameňov valley, 1450 m a. s. l., leg. Šmarda, April 1957, specimen Nr.1/00761
3. The Belanian Tatra Mts, Dolina Siedmich prameňov valley, limestone, 1350 m a. s. l., leg. Šmarda, August 1960, specimen Nr.1/00783
4. The High Tatra Mts, Jahňací štít peak, soil, 2000 m a. s. l., leg. Šoltés, September 20, 1980, specimen Nr.1/01720
5. The High Tatra Mts, Tatranská Lomnica, intravilán, granite rock, 838 m a. s. l., leg. Šoltés, November 24, 1978, specimen Nr.1/01763
6. The High Tatra Mts, Tatranská Lomnica, intravilán, granite rock, 839 m a. s. l., leg. Šoltés, October 28, 1979, specimen Nr.1/01765
7. The High Tatra Mts, Bielovodská dolina valley, limestone, *Neckeretum crispae*, 1120 m a. s. l., leg. Šoltés, August 9, 1983, specimen Nr.1/04040
8. The High Tatra Mts, Mengusovská dolina valley, under Kôpky peak, *Calamagrostietum villosae*, soil, 1680 m a. s. l., leg. Šeffler, July 26, 1982, specimen Nr.1/04323

9. The West Tatra Mts, Tichá dolina valley, on a touristic footpath, soil, 1380 m a. s. l., leg. Šoltés, October 15, 1984, specimen Nr.1/04740
10. The High Tatra Mts, Kačacia dolina valley, the bank of Kačacie pleso lake, granite rocks, 1590 m a. s. l., leg. Šoltés, August 21, 1984, specimen Nr.1/04783
11. The High Tatra Mts, Kriváň peak, above Jamské pleso lake, touristic footpath, soil, 1570 m a. s. l., leg. Šoltésová, November 1, 1984, specimen Nr.1/04879
12. The High Tatra Mts, Velická dolina valley, on a touristic footpath, soil, 1690 m a. s. l., leg. Šoltésová, July 26, 1984, specimen Nr.1/04955
13. The High Tatra Mts, Kôprová dolina valley, fork to Temnosmrečinová dolina valley, soil, 1680 m a. s. l., leg. Šoltésová, July 17, 1984, specimen Nr.1/05034
14. The High Tatra Mts, Kriváň peak, *Junco-Oreochloetum*, soil, 1980 m a. s. l., leg. Šoltésová, August 24, 1985, specimen Nr.1/05056
15. The Belanian Tatra Mts, Skalka hillock, sandy soil in a spruce forest, 900 m a. s. l., leg. Šoltés, October 12, 1986, specimen Nr.1/05351
16. The High Tatra Mts, Dlhý les kežmarský forest, rocky wall of a bridge, 790 m a. s. l., leg. Šoltés, June 4, 1993, specimen Nr.1/07269
17. The Belanian Tatra Mts, Monkova dolina valley, limestone, 1310 m a. s. l., leg. Šoltés, September 16, 1993, specimen Nr.1/07285
18. The Belanian Tatra Mts, Suchá dolina valley, calcic soil, 820 m a. s. l., leg. Šoltés, September 24, 1993, specimen Nr.1/07626
19. The High Tatra Mts, Dolina Zeleného plesa valley, *Luzuletum spadiceae*, soil, 1661 m a. s. l., leg. Šoltés, July 6, 2004, specimen Nr.1/15237
20. The High Tatra Mts, Lomnický štít peak, top area, *Oxyrio-Saxifragetum carpaticae*, 2630 m a. s. l., leg. Šoltés, August 11, 2004, specimen Nr.1/15323

There is no information on the overall occurrence of *Racomitrium elongatum* in Slovakia but we think it is as abundant as *Racomitrium canescens*. According to BEDNAREK-OCHYRA (1995), the moss is growing abundantly in dry, sunny, sandy or clay habitats, in pine forests, on steep path sides, on dunes, in heaths and on grassy places. In the Tatra Mts it occurs both on carbonate or granite bedrocks, from montane to the subalpine levels. The moss grows on sandy or humic soil, on rocks, bare soil, or directly on well trodden footpaths as well as covering old building structures. Its ecological requirements are similar to *Racomitrium canescens*, but requires more humid habitats.

The moss is part of the floristical composition of different communities, but doesn't show any relation to syntaxon (BEDNAREK-OCHYRA 1995). In the Tatra Mts, we have recorded *Racomitrium elongatum* in the communities of the alliances *Calamagrostion villosae* PAWŁOWSKI et al. 1928, *Festucion pictae* KRAJINA 1933, *Juncion trifidi* KRAJINA 1933, *Androsation alpinae* BR. – BL. in BR.-BL. et JENNY 1926 and in the moss community determined as *Neckeretum crispae* ŠMARDA 1947. Accompanied species on granite rocks include *Diplophyllum taxifolium* (WAHLENB.) DUMORT., *Gymnomitrium coralloides* NEES,

*Tritomaria exsecta* (SCHMIDEL ex SCHRAD.) SCHIFFNER ex LOESKE, *Andreaea rupestris* (SCHMIDEL ex SCHRAD.) SCHIFFNER ex LOESKE, *Grimmia incurva* SCHWÄGR., *Hydrogrimmia mollis* (BRUCH & SCHIMP.) LOESKE, *Kiaeria FALCATA* (HEDW.) I. HAGEN, *Racomitrium sudeticum* (FUNCK) BRUCH & SCHIMP., *Sanionia uncinata* (HEDW.) LOESKE and *Schistidium apocarpum* (HEDW.) BRUCH & SCHIMP. On acid soil associates include *Polytrichum juniperinum* HEDW., *Dicranella heteromalla* (HEDW.) SCHIMP., *Dicranum scoparium* HEDW., *Pleurozium schreberi* (WILLD. ex BRID.) MITT., *Pohlia nutans* (HEDW.) LINDB., *Hylocomium pyrenaicum* (SPRUCE) LINDB., *Hypnum cupressiforme* HEDW., *Sphagnum compactum* LAM. & DC., *Philonotis seriata* MITT., *Polytrichum alpinum* HEDW., *Pleurocladula albescens* (HOOK.) GROLLE, *Pellia neesiana* (GOTTSCHE) LIMPR., *Pohlia drummondii* (MÜLL. HAL.) A. L. ANDREWS, *Racomitrium lanuginosum* (HEDW.) BRID., *Oligotrichum hercynicum* (HEDW.) LAM. & DC., *Pohlia nutans* (HEDW.) LINDB. and *Pogonatum urnigerum* (HEDW.) P. BEAUV. On limestone substratum, the moss is associated with *Entodon concinnus* (DE NOT.) PARIS, *Schistidium strictum* (TURNER) LOESKE ex MARTENSSON, *Schistidium apocarpum* (HEDW.) BRUCH & SCHIMP., *Tortella tortuosa* (HEDW.) LIMPR., *Ditrichum flexicaule* (SCHWÄGR.) HAMPE, *Neckera crispa* HEDW., *Campyllum stellatum* (HEDW.) C.E.O. JENSEN, *Mnium thomsonii* SCHIMP., *Ptychodium plicatum* (SCHLEICH. ex F. WEBER & D. MOHR) SCHIMP., *Plagiochila porelloides* (TORR. ex NEES) LINDENB., *Leskea polycarpa* EHRH. ex HEDW.




Fig. 4. The distribution of *Racomitrium elongatum* in the Tatra Mts. Graphical source: TPN Zakopane

The species hasn't been evaluated against the old IUCN system in Slovakia (KUBINSKÁ *et al.* 2001), and since that time the species hasn't been included in the list of species occurring in Slovakia.

*Racomitrium elongatum* doesn't satisfy any criteria of the threatened categories of IUCN (ECCB 1995; HALLINGBÄCK 1998; HALLINGBÄCK 2001; HALLINGBÄCK *et al.*, 1998; IUCN 1994). In Czech Republic, the species is not redlisted as well (KUČERA & VÁŇA 2005).

## Acknowledgement

The project was supported by the VEGA, No 2/7070/27. The author is indebted to Peter Martin, Bristol, UK for correcting the English.

## References

- BEDNAREK-OCHYRA H. (1995): Rodzaj *Racomitrium* (*Musci*, *Grimmiaceae*) w Polsce: taksonomia, ekologia i fitogeografia. *Fragm. - Florist. Geobot., Series Polonica 2.*, 307 pp.
- DÜLL R. (1994): Deutschlands Moose. 2. Teil. IDH – Verlag, Bad Münstereifel, 211 pp.
- ECCB (European Committee for Conservation of Bryophytes). (1995): Red Data Book of European Bryophytes. Trondheim. 291 pp.
- FRISVOLL A. (1983): A taxonomic revision of the *Racomitrium canescens* group (*Bryophyta*, *Grimmiales*). - *Gunneria* 41: 1-181.
- HALLINGBÄCK T. (1998): The new IUCN categories tested on the Swedish bryophytes. - *Lindbergia* 23: 13-27.
- HALLINGBÄCK T. (2001): Globally Endangered Bryophyte Species in Europe. - *Novit. Bot. Univ. Carol.* 15: 9 – 26.
- HALLINGBÄCK T., HODGETTS N., RAEYMAEKERS G., SCHUMACKER R., SÉRGIO C., SÖDERSTRÖM L., STEWART N. & VÁŇA J. (1998): Guidelines for application of the revised IUCN threat categories to bryophytes. - *Lindbergia* 23: 6-12.
- IUCN (1994): IUCN Red List Categories. - IUCN, Gland.
- KUBINSKÁ A. & JANOVICOVÁ K. (1998): Machorasty, pp. 297-332. In MARHOLD, K., HINDÁK, F. (eds), *Zoznam nižších a vyšších rastlín Slovenska*, Veda, Bratislava.
- KUBINSKÁ A., JANOVICOVÁ K. & ŠOLTÉS R. (2001): Červený zoznam machorastov Slovenska, pp. 31-43. In BALÁŽ, D., MARHOLD, K., URBAN, P. (eds.), *Červený zoznam rastlín a živočíchov Slovenska*. - *Ochrana prírody* 20, Supplement.
- KUČERA J. & VÁŇA J. (2005): Seznam a červený seznam mechorostů České republiky (2005). - *Příroda*, Praha, 23: 1-104
- OCHYRA R. & BEDNAREK-OCHYRA H. (2004): A key to genera of the *Racomitrioideae*. - <http://www.mobot.org/plantscience/BFNA/V1/RacoConspectus.htm>
- VÁŇA J., KUČERA J., HRADÍLEK Z. & SOLDÁN Z. (2008): Mechorosty České republiky. - [http://bryoweb.prf.jcu.cz/klic/genera/racomitrium.html#Rac\\_elon](http://bryoweb.prf.jcu.cz/klic/genera/racomitrium.html#Rac_elon)

Received: November 11<sup>th</sup> 2008  
Revised: November 23<sup>rd</sup> 2008  
Accepted: November 23<sup>rd</sup> 2008

## ***Sorbus hugh-mcallisteri*, nom. nov. for *Sorbus apiculata* McALLISTER 2005, nom. illeg.**

VLASTIMIL MIKOLÁŠ

Hanojská 4, SK- 040 13 Košice, Slovakia, e-mail: [sorbusaria@azet.sk](mailto:sorbusaria@azet.sk)

---

Mikoláš V. (2008): *Sorbus hugh-mcallisteri* nom. nov. for *Sorbus apiculata* McAllister 2005, nom. illeg. – Thaiszia – J. Bot. 18: 65-67. – ISSN 1210-0420.

Abstract: *Sorbus apiculata* McALLISTER 2005 is a later homonym of *Sorbus apiculata* (KOVANDA) MIKOLÁŠ. The new name *Sorbus hugh-mcallisteri* MIKOLÁŠ is applied for the species. The species belongs to the section *Multijugae* (T.T.YU) McALL. It is known from NW Yunnan, China. *Sorbus apiculata* (KOVANDA) MIKOLÁŠ 2004 is a member of *Sorbus danubialis* agg. (*Sorbus* subg. *Aria*) with the occurrence in Bohemia, Czech Republic.

Keywords: *Sorbus*, *Sorbus apiculata*, *S. hugh-mcallisteri*, Czech Republic, China

---

HUGH McALLISTER, the best specialist of the genus of *Sorbus* s.s., recently described 28 species in his monograph of the genus (McALLISTER 2005). The author accepts the genus *Sorbus* in its strict sense (with the type species *S. acucuparia* L.), splitting it up into 2 subgenera: subgenus *Sorbus* and subgenus *Albocarmesinae* McALL. The latter includes species with fruits lacking chromoplasts and with white to crimson coloration. Some species have crimson petals, the character parallel to the European subg. *Chamaemespilus* (MEDICUS) K.KOCH of the genus *Sorbus* (or *Aria* (PERSOON) HOST if we accept the narrow concept of the genus *Sorbus* used by McALLISTER). The species of the subgenus are distributed in the Sino-Himalayan floristic region. *Sorbus apiculata* McALL. belongs to section *Multijugae* (T.T.YU) McALL., including numerous microspecies. Other sections embraced in the subgenus *Albocarmesinae* are the following: sect. *Discolores* (T.T.YU) McALL., *Insignes* (T.T.YU) McALL. (the correct name of this section is *Albocarmesinae*, because the type species *S. insignis* of the subgenus belongs here), *Hypoglaucae* McALL. and *Reductae*

[T.T.Yu) McALL. The section *Multijugae* is characteristic by combination of numerous leaflets (8 and more), soft fruits, flat inflorescence and petiole base not sheathing. *S. apiculata* McALL. (one of numerous microspecies of the section) is characterized by pearly white fruits, with moderately fleshy calyx lobes and 12-14 pairs of apiculate (name !), ovoid-oblong leaflets. It is one of probably apomictic microspecies in the section and its occurrence is recently known from NW part of Yunnan, China (NE of Zongdian, Hengduan Mts.), the well-known regional hot spot (KELLEY 2006).

*Sorbus apiculata* (KOVANDA) MIKOLÁŠ 2004 was validly published (comb. et stat. nov.) in the journal *Thaiszia* (vol. 13: p. 130, „2003“, 2004, basionym: *S. graeca* (SPACH) KOTSCHY var. *apiculata* KOVANDA, *Acta Dendrol. Českoslov.* 3: p. 60, *Icones*: p. 61, Fig. 11, 1961). This microspecies is the member of the subgenus *Aria*, close to *S. danubialis* (JÁV.) KÁRP. However, it is not probably a diploid species as it was published earlier (JANKUN et KOVANDA 1987). The species is probably tetraploid (VÍT 2006) and it is characteristic by a pointed apex of little, rhombic leaves. The microspecies is, perhaps, an endemic of Bohemia, Czech Republic. The type comes from the vicinity of Zbraslav, Hradiště hill (central Bohemia; today the locality belongs to the southernmost part of Prague). The microspecies is probably widely distributed in Bohemia (KOVANDA 1961, JANDA 2006, pers.comm.). KOVANDA (1961, p. 61) evaluates his *Sorbus graeca* var. *apiculata* as „not homogenous in its shut-in Bohemian area“. The typical specimens are (according to KOVANDA 1961) growing in phyllites of lower Vltava region. KOVANDA (1961) also gives data from the surroundings of Mohelno and Pavlov Hills (both in SE part of the Czech Republic) and even from surroundings of Hajnáčka and Humenné (central and eastern Slovakia). These plants, at least from the last locality (Mt. Sokol), are certainly not identical with this taxon (MIKOLÁŠ, pers. observ.). Probably *S. apiculata* (KOVANDA) MIKOLÁŠ is growing only in Bohemia. *S. danubialis* agg. includes also several other narrow microspecies in Bohemia (MIKOLÁŠ, pers. observ.; JANDA, pers. observ.).

Because the name *S. apiculata* (KOVANDA) MIKOLÁŠ was validly published in 2004, the use of name *S. apiculata* McALL. (McALLISTER H.: *The genus Sorbus. Mountain ash and other rowans*, p. 234: 51. *Sorbus apiculata* McALL., 2005) is to be considered as a later homonym of the former (taxonomically of different species), and it is necessary to substitute it. I hereby propose a nomen novum (replacement name) for it. The name underlines the basic importance of HUGH McALLISTER for taxonomy of *Sorbus* s.s.

***Sorbus hugh-mcallisteri* MIKOLÁŠ, nom. nov.** (*Sorbus apiculata* McALLISTER, *The genus Sorbus, The mountain ash and other rowans*, p. 234, nr. 51, The Royal Botanic Gardens, Kew, 2005), nom illeg., non *Sorbus apiculata* (KOVANDA) MIKOLÁŠ (*Thaiszia* 13: p. 130, „2003“, 2004).



## Acknowledgement

I am obliged for the first information on publication of the name *Sorbus apiculata* McALL. and data on the occurrence of *Sorbus apiculata* (KOVANDA) MIKOLÁŠ to Mr. P.JANDA (Slaný, Czech Republic) and RNDr. KAROL MARHOLD, CSc., senior lecturer (Bratislava) for nomenclatural advices and substantial English correction.

## References

- JANKUN A. et KOVANDA M. (1987): Apomixis and origin of *Sorbus bohemica* (Embryological studies in *Sorbus* 2). – *Preslia* 59: 97-116.
- KELLEY S.L. (2006): Resolving place names in AMDO & KHAM: A gazetteer for the Hengduan Mountains region of Southwest China. – *Acta Phytotax.Sin.* 44: 721-732.
- KOVANDA M. (1961): Taxonomical studies in *Sorbus* subg. *Aria*. – *Acta Dendrol.Čechoslov.* 3: 23-70.
- MCALLISTER H. (2005): The genus *Sorbus*. Mountain ash and other rowans. – The Royal Botanic Gardens, Kew.
- MIKOLÁŠ V. (2004): *Sorbus amici-petri* MIKOLÁŠ, a new hybridogenous species of the genus *Sorbus* s.l. from eastern Slovakia. – *Thaiszia* 13(„2003“): 127-133.
- VÍT P. (2006): Variabilita endemických zástupců rodu *Sorbus* L. v ČR: morfometrické, karyologické a molekulární zhodnocení (Variability of endemic members of the genus *Sorbus* L. in Czech Republic: morphometric, karyological and molecular evaluation). – Dipl. thesis (unpubl.).

Received: February 2<sup>nd</sup> 2008

Revised: December 1<sup>st</sup> 2008

Accepted: December 1<sup>st</sup> 2008



## Structure and adjective vegetations of grass-plots in Kyiv

OLGA MELEGHIK, VLADIMIR SOLOMAKHA<sup>1</sup> & TETYANA SOLOMAKHA<sup>2</sup>

<sup>1</sup>Taras Shevchenko's Kyiv University

<sup>2</sup>M.G. Kholodny Institute of Botany, National Academy of Sciences of Ukraine, Tereshchenkivska Street, 2, Kyiv-1, 01001, Ukraine, tel. (380-44) 234-5157, Fax: (380-44) 234-1064

---

Melegnik O., Solomakha V. & Solomakha T. (2008): Structure and adjective vegetations of grass-plots in Kyiv. – Thaiszia – J. Bot. 18: 69-74. – ISSN 1210-0420.

Abstract: The paper present of ecological floristic classification of Kyiv grass-plots. Ecological floristic classification of given 5 classes (*Molinio-Arrhenatheretea*, *Agropyretea repentis*, *Plantaginea majoris*, *Chenopodietea* and *Artemisietea vulgaris*), 8 alliances, 11 unions, 27 associations and 33 variants.

Floristic structure and qualitative adjectives of Kyiv grass-plots were investigated. Common evaluations of qualitative adjectives of Kyiv grass-plots coverings were cited. There are 63 alochtonic species among the ones found in the grass-plots on the territory of Kyiv. They include 35 species imported before the XVI century (archeophytes), and 28 brought later.

Keywords: grass-plots, syntaxonomia, vegetation, qualitative adjectives.

---

### Introduction

Cities native habitat is mainly represented as the man-made grass cover or diversely transformed natural one. Natural cover is the most sensitive component of urban landscapes. It is the first to perceive even marginal environment changes; respond on their degradation or disappearance of certain plants specimens. Therefore, grass-plots vegetation in cities requires constant care and regeneration.

Plant cover of urban territories is represented by the large number of species and is highly diverse. As a whole it could be grouped into three main decorative compositions:

- tree and bush plantations;
- flower compositions;
- grass-plots covers.

The contamination level of urban environment is dramatically increasing on the grounds of high scientific and technical progress pace, abrupt increase of newest technologies and industry intensification. As is known, vegetation is one of the most efficient stabilizing factors in the settlements. It is considered that optimum conditions for human being living in urban conglomerations are formed when trees and bushes make up by crown projection 20 - 30 % of explained by their universal peculiarities based on their permanent use independently of relief features and territory architecture.

Grass-plots perform important functions in urban environment. They improve environment's aestheticism, taking up the dust and toxic compounds, influencing on local ecological conditions. Another important characteristic of sod grasses in cities is their ability to hamper distribution of adventitious (quarantine, in particular) plants. Solid herbage and strong sod structure give no possibility to tap-root ruderal plants.

## **Material and methods**

Grass-plots in Kyiv of different functional purposes and on various stages of their development were studied. The experiments were conducted during 2002 – 2005 by the means of standard methods. All phytocoenoses were analysed in the field using methods of Zürich-Montpelliér school (BRAUN-BLANQUET 1964, WESTHOFF & VAN DER MAAREL 1978). Geobotanical descriptions were made according to the J. Braun-Blanquet method. Sample grounds were selected in different city regions. Their sizes have varied from 4 to 100 m<sup>2</sup>.

## **Results**

There are several grass-plots' classification schemes. Scheme designed by A. A. LAPTEV (1983) includes various in structure functional aspect grass-plots associations which are combined into more general categories.

According to the Mytsik L.P. classification there are five categories which completely cover all grass-plots characteristics: classic grass-plot, ordinary, cultural and natural.

We posed a task to determine complete floristic composition of grass-plots comparing to the classification scheme of vegetation of Ukraine. After handling of all descriptions the following classification scheme was obtained.

## Syntaxonomic vegetation scheme of grass-plots in Kyiv

- Cl. *Molinio-Arrhenatheretea* R.Tx. 1937
  - Ord. *Arrhenatheretalia elatioris* Pawl. 1928
    - Cynosurion cristati* Br.-Bl. et R.Tx. 1943
      - Ass. *Plantagini lanceolatae-Festucetum rubrae* Scamoni 1956
        - P.l.-F.r.* var. *Elytrigia repens*
        - P.l.-F.r.* var. *typicum*
        - Festucetum rubrae* Rübel 1912
          - F.r.* var. *Elytrigia repens*
          - F.r.* var. *typicum*
        - Lolio-Cynosuretum* Buker 1941
          - L.-C.c.* var. *Festuca rubra*
          - L.-C.c.* var. *Poa pratensis*
          - L.-C.c.* var. *typicum*
          - B.c.* *Lolium perenne* [*Cynosurion*]
        - Agrostio tenuis-Trifolietum repentis* (Walther 1977)
        - Leontodono-Poetum pratensis* Anishczenko et L. Ishb. 1989
          - L.-P.p.* var. *Elytrigia repens*
          - L.-P.p.* var. *Festuca rubra*
          - L.-P.p.* var. *typicum*
        - Poetum pratensis* Stepanovič 1999
          - P.p.* var. *Festuca rubra*
          - P.p.* var. *Trifolium repens*
          - P.p.* var. *Elytrigia repens*
          - P.p.* var. *typicum*
          - B.c.* *Poa pratensis* [*Cynosurion*]
- Agropyreteae repentis* Oberd., Th.Mull. et Gors in Oberd. et al. 1967
  - Agropyretalia repentis* Oberd., Th.Mull. et Gors in Oberd. et al. 1967
    - Convolvulo-Agropyron repentis* Gors 1966
      - Calamagrostietum epigeios* Kost. in V.Solomakha. et al. 1992
        - C. ep.* var. *typicum*
        - C. ep.* var. *Dactylis glomerata*
      - Convolvulo-Agropyretum repentis* Felf. (1942) 1943
        - C.-A.r.* var. *Potentilla argentea*
        - C.-A.r.* var. *Poa pratensis*
        - C.-A.r.* var. *typicum*
      - Agropyretum repentis* Gors 1966
        - A.r.* var. *Lolium perenne*
        - A.r.* var. *Poa pratensis*
        - A.r.* var. *typicum*
      - Poetum pratensis-compressae* Bornkamm 1974
- Polygono arenastri-Poëtea annua* Rivas-Martinez 1975 corr. Rivas-Martinez et al. 1991

*Plantaginietalia majoris* R. Tx. et Prsg. in R. Tx. 1950 em Rivas-Mart. 1975  
     *Polygonion avicularis* Br.-Bl. 1931  
         *Cynodonto-Plantaginietum majoris* Brun-Hool 1962  
         *Polygonetum avicularis* Gams 1927 em. Jehlik in Hejny et al. 1979  
         *Plantagini-Lolietum perennis* Beger 1930  
             *L.-P.m.* var. *Poa pratensis*  
             *L.-P.m.* var. *Trifolium repens*  
             *L.-P.m.* var. *typicum*  
         *Poetum annuae* Gams 1927  
             *P.a.* var. *Poa pratensis*  
             *P.a.* var. *typicum*  
         *Juncetum tenuis* (Diem., Siss. et Westh. 1940)  
         Schwick. 1944 em R. Tx. 1950  
         *Prunello-Plantaginietum majoris* Falinski 1963  
         *Agrostio tenuis-Poetum annuae* Gutte et Hilbig 1975  
*Agrostietalia stoloniferae* Oberd. in Oberd. et al. 1967  
     *Agropyro-Rumicion crispi* Nordhagen 1940  
         *Rumici crispi-Agrostietum stoloniferae* Moor 1958  
         *Potentillietum anserinae* Rap. 1927 em Poss. 1964  
*Chenopodietea* Br.-Bl. 1951 em Lohm., J. et R. Tx 1961 ex Matsz. 1962  
     *Sisymbrietalia* J. Tx. ex Matsz. 1962 em Gors. 1966  
         *Malvion neglectae* Gutte 1972  
             *Malvetum neglectae* Felf. 1942  
         *Sisymbrium officinalis* R. Tx., Lohm., Prsg. in R. Tx 1950 em Hejny et al. 1979  
         *Matricarietum perforatae* Kepczynska 1975  
             *M. p.* var. *Echinochloa crusgalli*  
             *M. p.* var. *typicum*  
         *Chenopodietum albi-viridae* Hejny 1979  
             *Ch. al.-v.* var. *typicum*  
             *Ch. al.-v.* var. *Lolium perenne*  
             *Ch. al.-v.* var. *Lolium multiflorum*  
             *Ch. al.-v.* var. *Portulaca oleracea*  
             *Ch. al.-v.* var. *Eragrostis minor*  
*Eragrostietalia* J. Tx. in Poli 1966  
     *Eragrostion* (R. Tx. 1950) Oberd. 1954  
         *Eragrostio-Amarantheum albi* Morariu 1943  
             *E.-A. al.* var. *Artemisia absinthium*  
             *E.-A. al.* var. *Festuca rubra*  
             *E.-A. al.* var. *Melilotus album*  
             *E.-A. al.* var. *Urtica dioica*  
             *E.-A. al.* var. *Typicum*

*Digitario-Portulacetum* (Felf. 1942) Timar et Bodrogkosi  
 1959  
     *D.-P. var. Berteroa incana*  
     *D.-P. var. Festuca ovina*  
     *D.-P. var. typicum*  
*Artemisietea vulgaris* Lohm., Prsg. et al. ex von Rochow 1951  
     *Meliloto-Artemisietalia absinthii* Elias 1979  
     *Potentillo-Artemision absinthii* Elias (1979) 1980  
     *Tanaceto-Artemisietum vulgaris* Br.-Bl. corr. 1949  
         *T.-A.v. var. typicum*  
         *T.-A.v. var. var. Lolium perenne*  
     *Dauco-Melilotion albi* Gors em Elias 1980  
         *Berteroetum incanae* Siss. et Tidem. ex Siss. 1950  
     *Artemisietalia vulgaris* Lohm. in R.Tx. 1947  
         *Arction lappae* R.Tx. 1937 em Gutte 1972  
         *Artemisietum vulgaris* R. Tx. 1942  
 D.c. *Lactuca serriola* [Sisimbrion officinalis]

As is well seen from the scheme sod covers' vegetation in Kyiv constitutes of 27 associations, 33 variants, 2 basal and 1 decorative communities, which belong to the 11 unions, 8 orders and 5 classes.

There are 63 alochtonic species among the ones found in the grass-plots on the territory of Kyiv. They include 35 species imported before the XVI century (archeophytes), and 28 brought later. The first group includes *Anisantha tectorum* (L.) Nevski, *Atriplex patula* L., *Lepidium ruderae* L., *Portulaca oleracea* L., *Ranunculus repens* L., *Setaria glauca* (L.) Beauv. etc., the second – *Ambrosia artemisiefolia* L., *Eragrostis minor* Host., *Lolium multiflorum* Lam., *Saponaria officinalis* L., *Trifolium hybridum* L., etc. The xenophytes/archeophytes correlation is 1/1.25 with archeophytes domination. Such ration is typical for Steppe zone of Ukraine and Polesie, while in southern regions xenophytes are prevailed (PROTOPOPOVA 1991).

By the importing means this species are divided into two groups: acolutophytes – species imported occasionally and the ones that are spread via anthropogenic broken ecotopes; and ergasiophytes – species imported in purpose for cultivation, and widespread into natural phytocenoses. Xenophytes are represented mainly by the adventive species: *Amaranthus albus* L., *Anagallis arvensis* L., *Bromus squarrosus* L., *Iva xanthifolia* Nutt., *Sisymbrium officinale* (L.) Scop. etc. Ergasiophytes are represented with the small number of species – *Trifolium hybridum*, *Saponaria officinalis* L., *Raphanus raphanistrum* L., *Lolium multiflorum*, *Lepidotheca suaveolens* etc.

By the naturalization degree in the natural flora research area two plant species groups are marked out – agriophytes (15) and epocophytes (48). Epocophytes are the species spread only in disturbed ecosystems. They form ruderal communities (*Amaranthus retroflexus* L., *Capsella bursa-pastoris* (L.) Medik., *Chenopodium polyspermum* L., *Descurainia sophia* (L.) Webb. et Prantl, *Digitaria sanguinalis* (L.) Scop., *Galinsoga parviflora* Cav., *Xanthoxalis stricta*

(L.) Small etc.). Epocophytes are resistant to the anthropogenic load which actually promotes their propagation and distribution. To agriophytes belong species widespread both in anthropogenic transformed landscapes and in natural phytocoenoses where they occupy a stable position. They include – *Juncus tenuis* Willd., *Phalacrolooma annuum*, *Saponaria officinalis*, *Ranunculus repens*, *Lamium maculatum* etc.

Correlation between indigenous and adventive fractions formed during the years is considered as very important characteristic of synantropic flora of every region since it indicates flora synantropization degree (PROTOPOPOVA 1991).

Now urban flora in Kyiv includes more than 600 adventive species which mainly are expansive. Plants causing mass allergic illnesses among the population (quarantine weeds) constitute a high menace for the human health – *Ambrosia artemisiifolia*, *Artemisia absinthium* L., *A. vulgaris* L., *Iva xanthifolia*, etc. Forming of solid and highly efficient sod cover on the cities territories prevents penetration and distribution of quarantine species; promotes environment optimization and decreases flora adventization degree in the region.

## Conclusions

Species composition of grass-plots in Kyiv includes 198 species of vascular plants, part of which are archeophytes (17, 68%) and part – kenophytes (14, 14%). Most of the allohtonic species are weeds, part of which is quarantine.

It was established that plant communities of grass-plots cultural phytocoenoses belong to the 5 vegetation classes, which combine 11 unions and 27 associations: *Molinio-Arrhenatheretea*, *Agropyretea repentis*, *Plantaginetea majoris*, *Chenopodietea* and *Artemisietea vulgaris*.

The ruderal degree of decorative covers of different functions was determined. The overall assessment of grass-plots quality indicates on low quality of cultural phytocoenoses in Kyiv.

## References

- BRAUN-BLANQUET J. (1964): Pflanzensoziologie. Grundzuge der Vegetationskunde. Ed. 3. – Springer-Verlag, Wien, New York. [865 pp.]
- LAPTEV A. A. (1983): Grass-plots. Kiev: Naukova Dumka Press – P. 3-106. [in Russ.].
- PROTOPOPOVA V. V. (1991): Synanthropic flora of Ukraine and ways of its development. Kiev: Naukova Dumka Press – 202 pp. [in Russ.].
- WESTHOFF V. & VAN DER MAAREL E. (1978): The Braun-Blanquet approach. – In: WHITTAKER R.H. (ed.): Classification of plant communities, Dr. W. Junk, The Hague, p. 287-399.

Received: November 06<sup>th</sup> 2008  
Revised: November 21<sup>st</sup> 2008  
Accepted: November 21<sup>st</sup> 2008



## Ecological analysis of the flora of saline sites in the northern part of Banat region in Vojvodina Province (Serbia)

ALEKSA KNEŽEVIĆ, SLOBODANKA STOJANOVIĆ, LJILJANA NIKOLIĆ, DEJANA DŽIGURSKI & BRANKA LJEVNAIĆ

Faculty of Agriculture, University of Novi Sad, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia, e-mail: ljnik@polj.ns.ac.yu

---

Knežević A., Stojanović S., Nikolić Lj., Džigurski D. & Ljevnaić B. (2008): Ecological analysis of the flora of saline sites in the northern part of Banat region in Vojvodina Province (Serbia). – Thaiszia – J. Bot. 18: 75-92. – ISSN 1210-0420.

Abstract: A long-term floristic study has determined the presence of 410 plant taxa in the flora of saline sites in northern Banat, the Vojvodina Province, Serbia. Of the determined taxa, 363 were included in the ecological analysis of the saline flora, i.e., all 348 determined species, all 14 determined subspecies and, because of its specific importance, only one variety, *Aster tripolium* var. *pannonicus* (Jacq.) Beck. Based on the analysis of mean values of ecological indexes (LANDOLT, 1977), we concluded that the saline soil of the studied region is moderately dry (F – 2.65), moderately basic (R – 3.30), moderately abundant with nitrogen (N – 3.08) and humus (H – 2.,95), poorly aerated (D – 3.89), highly saline (S<sub>+</sub> - 22.31 %), with a favorable thermal regime (T – 4.03), with the dominance of heliophytes (L – 3.85) which are well adapted to the conditions of the moderate continental climate (K – 3.24) that characterizes the studied region.

Keywords: Vojvodina Province, northern Banat, saline sites, flora, ecological index.

---

## Introduction

Most of the natural plant cover of the Vojvodina Province (Serbia) has been eradicated as the crop and animal productions gradually intensified. There still remains about 10% of land which has a fairly well-developed natural plant cover. These areas exhibit halophytic characteristics because those are predominantly salinas. In recent years these areas have been exposed to the anthropogenic factor at an increasing rate. It is characteristic for the plant cover of the saline sites in the Vojvodina Province that it had developed intrazonally within the zonal forest-steppe community of the alliances *Aceri tatarici-Quercion* Zólyomi et Jakucs 1957 and *Festucion rupicolae (sulcatae)* Soó (1940) 1964.

Numerous authors have studied the saline flora and vegetation (SLAVNIĆ 1948; BODROGKÖZY & GYÖRFFY 1970; PARABUĆSKI 1979, 1980; KUJUNDŽIĆ 1980; KNEŽEVIĆ 1994, BUDAK 1998, SÁDOVSKÝ et al., 2004; ELIÁŠ et al., 2008, DÍTĚ et al. 2008; KNEŽEVIĆ et al. 2008). In recent years, in addition to floristic and phytocoenological studies, we have focused our attention on the bioindication ecology of halophytes.

The use of ecological indices as indicator values provides an opportunity to characterize all aspects of a plant species in a succinct, simple and comparable way. When characteristics of plant species are defined in that manner, they may be used for the assessment of ecological characteristics and quality class of a given site. In other words, each plant species may serve as a test organism that reflects the conditions, status and quality of its environment. Such plant species are most sensitive biodiversity indicators which characterize not only parts of their environment but also the entire environment as an integral system, reflecting not only the consequences of processes that take place in the environment but also their causes (STOJANOVIĆ et al. 1994).

The objective of this study was to outline the ecological characteristics of saline sites in northern Banat, the Vojvodina Province, on the basis of an analysis of ecological indices of the recorded taxa.

## Materials and methods

The boundaries of the studied region are the watercourses of the rivers Tisza and Zlatica and the state border with Hungary and Romania. The saline sites in the region are predominated by limeless solonetz soils, which possess more or less columnar structure. Solonchak soils occur sporadically, in isolated spots. These soils had developed in consequence to the alluvial salinization by waters from the Carpathian basin (NEJGEBAUER et al. 1971). The subsequent regulation of the Tisza and Zlatica watercourses had prevented surface flooding, resulting in the occurrence of salt leaching in the studied saline sites.

The ecological indexes used in the analysis of basic environmental factors (humidity - F, chemical reaction - R, content of nitrogen and nitrogen compounds - N, content of humus (organomineral substances) - H, dispersion (aeration) of the substrate - D, salinity - S, light - L, temperature - T, and continentality - K) were taken from the publication 'Ökologische Zeigerwerte zur Schweizer Flora'

(LANDOLT 1977). The taxa which had not been mentioned by LANDOLT were characterized according to the publication "Monograph of Flora of Vascular Plants on the Saline Soils in the Banat Region (Yugoslavia)" (KNEŽEVIĆ 1994). In that publication, the criteria for determination of ecological indices were the papers of SOÓ (1964-1980), BORHIDY (1993), BODROGKÖZY (1970), KNEŽEVIĆ (1983), KNEŽEVIĆ et al. (1994, 1996, 1998, 2005, 2008), JANJATOVIĆ et al. (1990, 1992, 1995) and MERKULOV et al. (1999) and our own on-site observations. The supplement provides ecological indices for the recorded species, subspecies and one variety. The ecological indices supplied by LANDOLT are given in **bold** characters and those estimated by ourselves in normal characters.

## Results and discussion

In a long-term floristic study of the flora of saline sites in the examined region, presence of 410 taxa of vascular plant was determined (KNEŽEVIĆ et al. 2005). Of the determined taxa, 363 were included in the ecological analysis of the saline flora, i.e., all 348 determined species, all 14 determined subspecies and, because of its specific importance, only the variety *Aster tripolium* var. *pannonicus* (Jacq.) Beck.

Tab. 1 shows the following parameters for the analyzed taxa: number of taxa with a certain numerical value of ecological index, their percentages and, except for the relationship with salinity, mean values of the ecological indexes.

### Relationship between plants and humidity of the site (F)

The analysis of the ecological indexes for humidity (F) indicated the predominance of plant species with the ecological indexes  $F_2$  (131 taxa; 36.09%), adapted to dry sites, and  $F_3$  (102 taxa; 28.10%), adapted to moderately dry sites. The number of plant species with the ecological index  $F_1$  was lower (53 taxa; 14.60%). Those were xerophytes, i.e., indicators of extremely dry sites. After the xerophytes there followed the indicators of humid sites, i.e., the plant species with the ecological index  $F_4$  (45 taxa; 12.40%). The plant species with the ecological index  $F_5$  were fewest (32 taxa; 8.81%). This group contained indicators of high humidity of the site.

The mean value of the ecological indexes for humidity for the analyzed taxa was 2.65, indicating that the saline sites in the studied region are moderately dry.

### Relationship between plants and chemical reaction of the site (R)

There was a large dominance of neutrophilous species with the ecological index  $R_3$  (212 taxa; 58.40%). The proportion of indicators of neutral to basic chemical reaction of the site, designated with the ecological index  $R_4$ , was also significant (129 taxa; 35.54%). There was a small number of indicators of acidic substrate, with by the ecological index  $R_2$  (20 taxa; 5.51%). The presence of indicators of highly acidic sites, labeled as  $R_1$ , and indicators of highly basic sites, labeled as  $R_5$ , was minimal (1 taxon each; 0.27%).

The mean value of the ecological indexes for chemical reaction of the site was 3.30, indicating that the saline sites in the studied region are moderately basic.

**Tab. 1. Ecological analysis of flora of saline sites in the northern part of Banat region in the Vojvodina Province**

<b>Ecological index</b>	<b>Numerical values of ecological index</b>	<b>Number of taxa</b>	<b>%</b>	<b>Mean value</b>
<b>F – humidity</b>	1	53	14.60	<b>F – 2.65</b>
	2	131	36.09	
	3	102	28.10	
	4	45	12.40	
	5	32	8.81	
<b>R – chemical reaction of the site</b>	1	1	0.27	<b>R - 3.30</b>
	2	20	5.51	
	3	212	58.40	
	4	129	35.54	
	5	1	0.27	
<b>N - nitrogen and nitrogen compounds</b>	1	21	5.78	<b>N - 3.08</b>
	2	81	22.31	
	3	124	34.16	
	4	120	33.06	
	5	17	4.68	
<b>H – humus</b>	1	2	0.55	<b>H - 2.95</b>
	2	61	16.80	
	3	255	70.25	
	4	42	11.57	
	5	3	0.83	
<b>D – dispersion (aeration)</b>	1	0	0	<b>D - 3.89</b>
	2	5	1.38	
	3	119	32.78	
	4	149	41.05	
	5	90	24.79	
<b>S – salinity</b>	-	282	77.69	
	+	81	22.31	
<b>L – light</b>	1	0	0	<b>L - 3.85</b>
	2	2	0.55	
	3	82	22.59	
	4	264	72.73	
	5	15	4.13	
<b>T – temperature</b>	1	0	0	<b>T – 4.03</b>
	2	1	0.27	
	3	72	19.83	
	4	204	56.20	
	5	86	23.69	
<b>K – continentality</b>	1	1	0.27	<b>K - 3.24</b>
	2	45	12.40	
	3	195	53.72	
	4	108	29.75	
	5	14	3.86	

### **Relationship between plants and nitrogen and nitrogen compounds (N)**

The analysis of the ecological indexes for the contents of nitrogen and nitrogen compounds indicated the prevalence of taxa with the ecological indexes  $N_3$  (124 taxa; 34.16%) and  $N_4$  (120 taxa; 33.06%). The proportion of taxa characterized by the index  $N_2$  was significantly lower (81 taxa; 22.31%). The numbers of indicators of very poor sites, designated with the index  $N_1$ , and indicators of sites rich in nutrients, designated with the index  $N_5$ , were low (21 taxa or 5.78% and 17 taxa of 4.68%, respectively).

Based on the mean value amounting to 3.08, the saline sites in the studied region were considered to be moderately rich in biogenous nutrients.

### **Relationship between plants and humus content (H)**

The analysis of the ecological indexes for the content of humus indicated a high predominance of taxa with the ecological index  $H_3$  (255 taxa; 70.25%). This characterizes the studied saline sites as moderately rich in organomineral compounds. The indicators of low humus content designated with the ecological index  $H_2$  were considerably fewer (61 taxa; 16.80%). The relative abundance of humus in the substrate was indicated by a small number of taxa with the ecological index  $H_4$  (42 taxa; 11.57%). The numbers of indicators of humic sites, designated with the ecological index  $H_5$ , and indicators of sites poor in humus, designated with the ecological index  $H_1$ , were quite low (3 taxa or 0.83% and 2 taxa or 0.55%, respectively).

The mean value of the ecological index for humus content was 2.95, describing the studied site as moderately provided with humus.

### **Relationship between plants and dispersion (aeration) of the substrate (D)**

Almost two thirds of the analyzed taxa were designated with the ecological indexes  $D_4$  (149 taxa; 41.05%) and  $D_5$  (90 taxa; 24.79%). Those were plant species adapted to a non-skeletal soil with a heavy mechanical composition. A significant presence of plant species designated with the ecological index  $D_3$  (119 taxa; 32.78%) testified that the saline sites included also some well aerated locations. Indicators of soil with large particles, designated with the ecological index  $D_2$ , were scarce (5 taxa; 1.38%), while indicators of immature soil, i.e., plant species labeled with the ecological index  $D_5$ , could not be found at the saline sites of the studied region.

The mean value amounting to 3.89 indicates that the studied saline soils had unfavorable dispersion, i.e., poor aeration.

### **Relationship between plants and salinity (S)**

The taxa designated with the ecological index  $S_+$  made more than one fifth of the analyzed flora (81 taxa; 22.31%). This gives the halophytic character to the flora, despite the dominance of plants species designated with the ecological index  $S$ . (282 taxa; 77.69%). The previous statement is based on the fact that the taxa bearing the ecological index  $S$  include numerous plants species which are tolerant to salinity as well as the species which avoid high salinity of the site by

completing their vegetation in the spring, when salt concentration in the substrate is low due to abundant rainfall.

### **Relationship between plants and light (L)**

Plant species designated with the ecological index  $L_4$  were dominant in the analyzed flora (264 taxa; 72.73%). These species are adapted to full light but they also tolerate a certain degree of shading. The number of taxa with the ecological index  $L_3$  was considerably lower (82 taxa; 22.59%). These species are indicators of half-shade. There was a small number of species with the ecological index  $L_5$  (15 taxa, 4.13%). These species do not tolerate shading. The species designated with the ecological index  $L_2$  were few (2 taxa, 0.55%). They are indicators of full shade. Indicators of deep shade, designated with the ecological index  $L_1$ , were absent.

The mean value of 3.85 was an indication of dominance of heliophytes ( $L_4$ ) over semi-sciophytes ( $L_3$ ).

### **Relationship between plants and temperature (T)**

The analysis of ecological indexes for temperature showed that thermophilous plant species designated as  $T_4$  (204 taxa, 56.20%) and true thermophiles designated as  $T_5$  (86 taxa, 23.69%) dominated the flora of the studied saline sites. The participation of taxa with the ecological index  $T_3$  was lower (72 taxa, 19.83%). These species were indicators of moderately warm sites. The presence of taxa designated as  $T_2$ , indicators of cold sites, was insignificant (1 taxon, 0.27%). Taxa from very cold sites, designated as  $T_1$ , were completely absent.

The mean value of 4.03 indicated a favorable temperature regime of the sites.

The analysis of adaptation of the plant species of the studied flora to the continental climate showed that the largest number of plant species were designated with the ecological index  $K_3$  (195 taxa, 53.72%). This was an indication that moderately continental conditions dominated in the studied region. A considerable influence of typical continental conditions resulted in a large presence of taxa designated with the ecological index  $K_4$  (108 taxa, 29.75%). Plant species with the ecological index  $K_2$ , indicating an influence of sub-oceanic climate, were present in lower numbers (45 taxa, 12.40%). The number of plant species of typical continental regions, designated with the ecological index  $K_5$ , was small (14 taxa, 3.86%). The presence of oceanic plant species, designated with the ecological index  $K_1$ , was insignificant (1 taxon, 0.27%).

The mean value of 3.24 confirmed that the moderate continental climate of the studied region was under a considerable influence of the typical continental climate.

Based on the bioindicator values of the analyzed flora, the saline sites in the region surrounded by the rivers Tisza and Zlatica and the state border with Hungary and Romania may be characterized as moderately dry, moderately basic, moderately abundant with nutrients, moderately abundant with humus, but poorly aerated. Their plant cover has halophytic characteristics, with heliophytes predominating under a favorable thermal regime of the moderate continental climate of the studied region.

The obtained results indicate that the saline pastures of northern Banat have a low annual biomass production, as observed earlier for the saline pastures of southeastern Banat (VUČKOVIĆ 1985). Their more productive parts are used primarily for grazing and hay production, and only exceptionally for crop production. The poorest sites should be used for the construction of fishponds and infrastructure facilities, in order to preserve the neighboring tracts of fertile land. Precise locations for the construction of these facilities could be recommended by analyzing the indicator values of individual stands of the studied plant communities.

## Acknowledgements

This study is part of the project TR – 20083 titled »Improvement of technology of production of forage crops in the function of the production of safe cattle feed«, subsidized by the Ministry for Science and Technological Development of the Republic of Serbia.

## References

- BODROGKÖZY GY. & GYÖRFFY B. (1970): Ecology of the halophytic vegetation of the pannonicum VII. Zonation study along the Bega-backwaters in the Voivodina (Yugoslavia). Acta Biologica Hungarica. Tom XVI (3-4), Szeged.
- BORHIDI A. (1993): A Magyar Flóra szociális magatartás típusai, természetessége és relativ ökológiai értékszámai. Janus Pannonius Tudományegyetem, Pécs, p. 95.
- BUDAK V. (1998): Flora i biljnogeografske odlike flore slatina Bačke. Matica srpska. Novi Sad, p. 191.
- DÍTĚ D., ELIÁŠ P. & SÁDOVSKÝ M. (2008): *Camphorosmetum annuae* Rapaics ex Soó 1933 – vanishing plant community of saline habitats in Slovakia. Thaiszia – J. Bot. Košice, 18:9-20.
- ELIÁŠ P., DÍTĚ D., GRULICH V. & SÁDOVSKÝ M. (2008): Distribution and communities of *Crypsis aculeata* and *Heleochoa schoenoides* in Slovakia. Hacquetia 7/1: 5-20.
- JANJATOVIĆ V., KNEŽEVIĆ A. & KABIĆ D. (1990): Ekomorfološka istraživanja *Aster tripolium* L. var. *pannonicus* (Jacq.) Beck. Bilten društva ekologa Bosne i Hercegovine, ser B, br.5 71 -75, Sarajevo.
- JANJATOVIĆ V., KNEŽEVIĆ A., ANĐELIĆ M. & MERKULOV L.J. (1992): Eko-morfološke karakteristike vrste *Camphorosma annua* Pall. (*Chenopodiaceae*). Zbornik radova PMF, ser. biol. 22, 31-38. Novi Sad.
- JANJATOVIĆ V., MERKULOV L.J. & KNEŽEVIĆ A. (1995): Prilog proučavanju ekologije *Atriplex litoralis* L. (*Chenopodiaceae*). Zbornik radova PMF, ser. biol. 24, 51 - 57, Novi Sad.
- KNEŽEVIĆ A. (1983): Prilog proučavanju halofitske vegetacije severnog Banata. Zbornik za prirodne nauke Matice srpske, 65, 83-94, Novi Sad.
- KNEŽEVIĆ A. (1994): Monografija flore vaskularnih biljaka na slatinama u regionu Banata (Jugoslavija). Matica srpska, Novi Sad, p. 122 (in Serbian, with English abstract)
- KNEŽEVIĆ A., BUTORAC B. & BOŽA P. (1994): Ecological and phytogeographical analysis of vegetation of the alliance *Halo-Agrostion albae pannonicum* Knežević 1990 (syn. *Beckmannion eruciformis* Soó 1933). Glasnik Instituta za botaniku i Botaničke bašte Univerziteta u Beogradu, 28, 159-172. Beograd.

- KNEŽEVIĆ A., MERKULOV L.J., BOŽA P. & GRDINIĆ B. (1996): Eko-morfološke adaptacije vrste *Roripa kernerii* Menyh. 1877 (*Caprales, Brassicaceae*). Zbornik radova PMF, ser. biol. 25, 29-36, Novi Sad.
- KNEŽEVIĆ A., MERKULOV L.J. & BOŽA P. (1998): Eco-morphological adaptations of *Puccinellia limosa* (Schur) Holmb. (*Poales, Poaceae*). Zbornik Matice srpske za prirodne nauke, br. 95, 59-69, Novi Sad.
- KNEŽEVIĆ A., STOJANOVIĆ S., BOŽA P., LAZIĆ D. & NIKOLIĆ L.J. (2005): Floristic composition and vegetation cover of saline sites in the northern part of Banat region in the Vojvodina Province. 8<sup>th</sup> International Symposium Interdisciplinary Regional Research Hungary - Serbia and Montenegro, Szeged, 2005, Proceedings, EEP, pp.19.
- KNEŽEVIĆ A., LJEVNAIĆ B., NIKOLIĆ L.J. & ČUPINA B. (2008): Biljnogeografska analiza flore zaslanjenih pašnjaka severnog dela vojvođanskog Banata. Acta Biologica Jugoslavica, serija G: Acta herbologica, 17 (1): 105-108.
- KUJUNDŽIĆ M. (1980): Slatinska vegetacija u okolini Ruskog Krstura (Vegetation of Saline Soils Near Ruski Krstur). Zbornik Matice srpske za prirodne nauke, Novi Sad, No.58: 99-112.
- LANDOLT E. (1977): Ökologische Zeigerwerte zur Schweizer Flora, Veröffentlichungen des Geobotanischen Institutes der ETH, Stiftung Rübel, 64. Heft. Zürich.
- MERKULOV L.J., KNEŽEVIĆ A. & BOŽA P. (1999): A comparative anatomical analysis of the species of genera *Crypsis* Ait. 1789. and *Heleochoa* Host 1801. (*Poales, Poaceae*). Zbornik Matice srpske za prirodne nauke, br. 96, 41-50, Novi Sad.
- NEJGEBAUER V., ŽIVKOVIĆ B., TANASJEVIĆ Đ. & MILJKOVIĆ N. (1971): Pedološka karta Vojvodine. Institut za poljoprivredna istraživanja. Novi Sad.
- PARABUĆSKI S. (1979): Zajednice *Peucedano-Asteretum punctati* Soó i *Trifolietum subterranei* Slavnić na nekim lokalitetima Bačke i njihov sintaksonomski položaj. Zbornik Matice srpske za prirodne nauke, Novi Sad, No.56: 17-43.
- PARABUĆSKI S. (1980): Karakteristike nekih halofitskih fitocenoza u Bačkoj. Zbornik Matice srpske za prirodne nauke, Novi Sad, No.58: 81-98.
- SÁDOVSKÝ M., ELIÁŠ P. & DITĚ D. (2004): Distribution of halophytic communities in southwestern Slovakia: history and present. Bull. Slov. Bot. Spoločn., Bratislava, Supl. 10:127-129.
- SLAVNIĆ Ž. (1948): Slatinska vegetacija Vojvodine. arhiv za biološke nauke i tehniku, Beograd, 3(4): 76-142.
- Soó R. (1964-1980): A magyar flóra és vegetáció Rendszeretani novényfoldrajzi kézikönyve I-VI, Akadémiai kiadó, Budapest.
- STOJANOVIĆ S., BUTORAC B., VUČKOVIĆ M., STANKOVIĆ Ž., ŽDERIĆ M., KILIBARDA P. & RADAK (NIKOLIĆ) L.J. (1994): Biljni svet kanala Vrbas-Bezdan. Univerzitet u Novom Sadu, Prirodno-matematički fakultet, Institut za biologiju, 1-110, Novi Sad (in Serbian, with English abstract)
- VUČKOVIĆ R. (1985): Fitocenoze slatinske vegetacije istočnog Potamišja, njihova produkcija i hranljiva vrednost. Doktorska disertacija odbranjena na PMF u Beogradu pp. 1- 217.



## Supplement

Taxon	Ecological index									
	F	R	N	H	D	S	L	T	K	
<i>Abutilon theophrasti</i> Meddik.	2	3	4	3	4	-	4	5	3	
<i>Achillea millefolium</i> L.	3	3	4	3	4	-	4	3	3	
<i>A. millefolium</i> L. subsp. <i>collina</i> (Becker) Weis.	2	4	2	3	3	-	4	4	4	
<i>A. millefolium</i> L. subsp. <i>pannonica</i> (Scheele) Hayek	2	3	4	4	3	-	4	4	4	
<i>A. setacea</i> W. et K.	1	3	2	2	3	-	4	5	5	
<i>Acorellus pannonicus</i> (Jacq.) Palla	4	4	4	2	4	+	4	4	4	
<i>Adonis aestivalis</i> L.	2	4	2	3	4	-	3	5	3	
<i>A. flammea</i> Jacq.	2	4	2	3	4	-	3	5	3	
<i>A. aestivalis</i> L.	2	4	2	3	4	-	3	5	3	
<i>Agrimonia eupatoria</i> L.	2	4	3	3	4	-	4	4	3	
<i>Agropyrum cristatum</i> (Schreb.) P.B.	2	3	2	3	3	-	4	4	4	
<i>A. repens</i> (L.) Beauv.	3	3	4	2	3	+	4	3	3	
<i>Agrostis alba</i> L.	4	3	3	3	4	-	4	5	2	
<i>Ajuga genevensis</i> L.	2	3	3	3	4	-	3	4	4	
<i>Alisma lanceolatum</i> With.	5	3	3	3	5	-	4	4	3	
<i>A. plantago-aquatica</i> L.	5	3	3	3	5	-	4	4	3	
<i>Allium atropurpureum</i> Waldst. Et Kit.	2	3	2	3	4	-	4	4	3	
<i>A. scorodoprasum</i> L.	4	3	4	4	5	-	3	5	3	
<i>A. sphaerocephalum</i> L.	1	4	2	2	3	-	4	4	4	
<i>A. vineale</i> L.	2	3	3	3	4	-	4	4	2	
<i>Alopecurus geniculatus</i> L.	4	3	4	3	5	+	3	4	3	
<i>A. pratensis</i> L.	4	3	4	3	5	-	4	3	3	
<i>Althaea officinalis</i> L.	3	3	4	3	4	-	4	5	4	
<i>Amarantus albus</i> L.	1	3	4	3	3	-	4	4	3	
<i>A. crispus</i> (Lesp. et Thev.) Terr.	2	3	4	4	3	-	4	4	5	
<i>A. retroflexus</i> L.	2	3	4	3	3	-	4	4	3	
<i>Anagallis arvensis</i> L.	3	3	3	3	4	-	4	4	3	
<i>Andropogon ischaemum</i> L.	2	3	2	2	3	-	5	5	4	
<i>Antriscus caucalis</i> M. B.	3	3	3	3	4	-	3	3	3	
<i>Arenaria serpyllifolia</i> L.	2	3	3	3	3	-	4	4	3	
<i>Arrhenatherum elatius</i> (L.) Mert. et Koch	3	3	4	3	4	-	3	4	3	
<i>Artemisia absinthium</i> L.	2	4	4	3	4	-	4	4	4	
<i>A. austriaca</i> Jacq.	2	3	3	3	3	-	4	4	4	
<i>A. maritima</i> L. subsp. <i>monogyna</i> (W. et K.) Gams.	2	4	2	2	4	+	4	4	5	
<i>A. maritima</i> L. subsp. <i>salina</i> (Willd.) Gams.	2	4	2	2	4	+	4	4	5	
<i>A. vulgaris</i> L.	3	3	4	3	4	-	4	4	3	
<i>Aster tripolium</i> L. var. <i>pannonicus</i> (Jacq.) Beck	3	4	2	2	4	+	4	4	4	
<i>Astragalus cicer</i> L.	2	4	4	3	3	-	3	4	4	
<i>Atriplex hastata</i> L.	3	3	4	3	3	+	4	4	4	
<i>A. litoralis</i> L.	3	3	2	2	4	+	3	4	3	
<i>A. tatarica</i> L.	3	3	4	2	4	+	3	3	4	
<i>Ballota nigra</i> L.	2	3	5	3	4	-	4	5	4	
<i>Beckmannia eruciformis</i> (L.) Host.	5	4	4	3	5	+	4	3	4	

Supplement – cont.

Taxon	Ecological index									
	F	R	N	H	D	S	L	T	K	
<i>Bellis perennis</i> L.	3	3	4	3	4	-	4	3	3	
<i>Berteroa incana</i> (L.) DC.	1	3	3	3	3	-	4	4	4	
<i>Bidens cernuus</i> L.	5	3	5	4	4	-	4	4	3	
<i>B. tripartitus</i> L.	5	3	5	4	4	-	4	4	3	
<i>Bolboschoenus maritimus</i> (L.) Palla	5	4	3	3	5	+	4	4	3	
<i>B. maritimus</i> (L.) Palla var. <i>compactus</i> (Hoffm.)										
<i>B. maritimus</i> (L.) Palla var. <i>macrostachys</i> (Wild) Kneuc. f. <i>macrostachys</i>										
<i>Bromus comutatus</i> Schrad.	2	3	3	3	5	-	3	4	3	
<i>Bromus comutatus</i> Schrad. f. <i>violaceus</i> Podp.										
<i>Bromus erectus</i> Huds.	2	4	2	3	4	-	4	4	3	
<i>B. inermis</i> Leyss.	2	4	3	3	3	-	4	4	4	
<i>B. mollis</i> L.	3	3	4	3	4	-	4	4	3	
<i>Bupleurum pachnospermum</i> Pančić	2	3	3	4	3	-	4	5	3	
<i>B. tenuissimum</i> L.	3	4	3	3	5	+	4	5	1	
<i>Butomus umbellatus</i> L.	5	3	4	4	5	-	3	4	3	
<i>Calystegia sepium</i> (L.) R. Br.	4	4	4	3	5	-	3	4	2	
<i>Camphorosma annua</i> Pall.	2	4	1	1	4	+	5	4	4	
<i>C. annua</i> Pall. f. <i>nana</i> Moq.										
<i>Capsella bursa pastoris</i> (L.) Medik.	2	3	4	3	4	-	4	3	3	
<i>Carduus acanthoides</i> L.	1	3	4	3	3	-	4	5	2	
<i>C. nutans</i> L.	2	4	4	2	3	-	4	4	3	
<i>Carex divisa</i> Huds.	4	4	3	3	3	+	5	5	2	
<i>C. hirta</i> L.	3	3	3	3	4	-	3	4	3	
<i>C. nutans</i> Huds.	3	3	3	3	3	-	4	5	4	
<i>C. praecox</i> Schreb	2	4	2	2	3	-	4	4	4	
<i>C. pseudocyperus</i> L.	5	4	3	4	5	-	4	4	4	
<i>C. spicata</i> Huds.	3	3	4	3	5	-	4	3	3	
<i>C. stenophylla</i> Wahlbg.	2	4	2	3	3	-	4	5	5	
<i>C. vulpina</i> L.	5	4	2	4	4	-	3	4	3	
<i>C. vulpina</i> L. f. <i>minor</i> Peterm.										
<i>C. vulpina</i> L. f. <i>longibracteata</i> Beck										
<i>Carthamus lanatus</i> L.	1	3	4	3	3	-	4	5	3	
<i>Centaurea calcitrapa</i> L.	2	4	4	3	5	-	4	5	2	
<i>C. cyanus</i> L.	2	3	3	3	4	-	4	4	4	
<i>C. difusa</i> Lam.	1	4	4	2	3	-	4	5	3	
<i>C. jacea</i> L. subsp. <i>angustifolia</i> (Schrk.) Gugl.	3	3	2	4	4	-	4	4	3	
<i>C. jacea</i> L. subsp. <i>angustifolia</i> (Schrk.) Gugl. var. <i>pannonica</i> (Heuff.) Gugl.										
<i>C. scabiosa</i> L. subsp. <i>sadleriana</i> (Janka) Aschers. et Graebn.	2	4	2	2	3	-	4	4	4	
<i>C. solstitialis</i> L.	2	3	4	2	4	-	4	4	4	
<i>Centaureum pulchellum</i> (Sw.) Druce.	3	4	3	2	5	+	4	4	3	
<i>C. uliginosum</i> (W. et K.) Beck.	3	4	1	3	4	+	4	4	5	

Supplement – cont.

Taxon	Ecological index								
	F	R	N	H	D	S	L	T	K
<i>C. umbellatum</i> Gilib.	3	3	3	3	4	-	4	4	3
<i>C. umbellatum</i> Gilib. f. <i>humile</i> (Dvorak) Soó									
<i>Cephalaria transsilvanica</i> (L.) Schrad.	1	4	1	1	2	-	4	4	3
<i>Cephalaria transsilvanica</i> (L.) Schrad. var. <i>allionii</i> (Kern.) Ravy									
<i>Cerastium arvense</i> L.	2	3	3	3	4	-	4	4	3
<i>C. banaticum</i> (Roch.) Heuff.	3	3	3	3	4	-	4	3	3
<i>C. brachypetalum</i> Desp.	1	4	1	3	3	-	4	4	3
<i>C. caespitosum</i> Gilib.	3	3	3	3	4	-	3	3	3
<i>C. dubium</i> (Bast.) Schwarz.	4	3	3	3	4	-	3	5	2
<i>C. pumilum</i> Curt.	1	4	2	3	3	-	4	4	2
<i>C. semidecandrum</i> L.	2	4	2	3	3	-	4	4	3
<i>Chenopodium album</i> L.	2	3	4	3	4	-	4	3	3
<i>Ch. glaucum</i> L.	3	3	5	3	4	+	4	4	4
<i>Ch. polyspermum</i> L.	3	3	4	3	4	-	4	4	3
<i>Ch. rubrum</i> L.	3	3	5	3	4	+	4	3	4
<i>Ch. rubrum</i> L. subsp. <i>botryo</i> (i) des Sm.	3	3	5	3	4	+	4	3	4
<i>Ch. vulvaria</i> L.	2	3	5	3	3	-	4	4	4
<i>Chondrilla juncea</i> L.	2	3	3	3	3	-	4	5	3
<i>Chrysopogon gryllus</i> (L.) Trin.	1	3	2	2	3	-	4	5	3
<i>Ch. gryllus</i> (L.) Trin. f. <i>flavencens</i> (Schur) Soó									
<i>Cichorium intybus</i> L.	2	4	3	3	5	-	5	4	3
<i>Cirsium arvense</i> (L.) Scop.	3	3	4	3	4	+	3	4	3
<i>C. arvense</i> (L.) Scop. var. <i>vestitum</i> Wimm. et Grab.									
<i>C. canum</i> (L.) All.	4	3	2	3	3	-	4	3	4
<i>C. lanceolatum</i> (L.) Scop.	3	3	4	4	4	-	3	3	3
<i>Consolida orientalis</i> (Gay.) Schröd.	2	3	4	3	3	-	3	4	2
<i>C. regalis</i> S.F.Gray.	2	5	3	3	3	-	3	4	4
<i>Convolvulus arvensis</i> L.	2	4	3	3	4	-	4	4	3
<i>C. arvensis</i> L. f. <i>microphyllus</i> Opiz									
<i>Corispermum nitidum</i> Kit.	2	4	2	3	3	-	4	4	4
<i>Crepis capillaris</i> (L.) Wallr.	3	3	3	3	4	-	4	4	3
<i>C. foetida</i> L. subsp. <i>rhoadifolia</i> (M.B.) Fiori et Paol.	1	3	4	3	3	-	4	5	3
<i>C. setosa</i> Hall.	1	3	4	3	3	-	4	5	3
<i>C. setosa</i> Hall. f. <i>gracilis</i> Rohlena									
<i>Crypsis aculeata</i> (L.) Aitt.	3	4	1	2	5	+	4	4	4
<i>Cynodon dactylon</i> (L.) Pers.	2	3	3	3	3	-	4	5	2
<i>Daucus carota</i> L.	2	3	2	3	3	-	4	4	3
<i>Descurainia sophia</i> (L.) Weeb.	2	3	4	3	4	-	4	3	4
<i>D. sophia</i> (L.) Weeb. var. <i>sophia</i> f. <i>minus</i> Bolzon ap. Hegi									
<i>Dianthus pontederæ</i> Kerner	2	4	3	3	3	-	4	4	4
<i>Dipsacus laciniatus</i> L.	3	4	4	3	5	-	4	5	3
<i>Echium vulgare</i> L.	1	3	4	2	3	-	5	4	4
<i>Epilobium adnatum</i> Griseb.	4	3	3	4	5	-	3	4	2

Supplement – cont.

Taxon	Ecological index									
	F	R	N	H	D	S	L	T	K	
<i>Eragrostis megastachya</i> (Koel.) Link	1	4	2	2	3	-	4	5	3	
<i>E. pilosa</i> (L.) P.B.	1	2	3	2	3	-	4	5	3	
<i>Erigeron canadensis</i> L.	2	3	3	3	4	-	4	4	3	
<i>Erodium cicutarium</i> (L.) L. Hérit.	2	3	3	3	3	-	4	4	3	
<i>Erophila verna</i> (L.) Schevall.	2	3	2	3	3	-	4	4	3	
<i>Eryngium campestre</i> L.	2	4	3	3	3	-	4	5	3	
<i>Euclidium syriaca</i> (L.) R. Br.	1	3	4	3	4	-	4	4	4	
<i>Euphorbia cyparissias</i> L.	2	3	2	3	4	-	4	3	3	
<i>E. glareosa</i> M.B.	1	4	1	4	4	-	4	4	4	
<i>E. palustris</i> L.	4	4	4	4	5	+	3	4	2	
<i>E. platyphyllos</i> L.	3	3	3	3	4	-	3	4	2	
<i>E. seguieriana</i> Neck.	1	4	2	3	2	-	4	4	4	
<i>Falcaria vulgaris</i> Bernh.	2	4	2	3	3	-	4	5	4	
<i>Festuca arundinacea</i> Schreb.	4	4	4	3	5	-	4	3	3	
<i>F. pratensis</i> Huds.	3	3	4	4	4	-	4	3	3	
<i>F. vallesiaca</i> Sch.	1	3	2	2	3	-	4	4	5	
<i>F. vallesiaca</i> Sch. subsp. <i>pseudovina</i> (Hack.) A. et G.	1	3	2	2	3	+	4	4	5	
<i>F. vallesiaca</i> Sch. subsp. <i>pseudovina</i> (Hack.) A. et G. f. <i>rutila</i> Hack.										
<i>Filago germanica</i> L.	1	2	3	2	3	-	4	5	3	
<i>Filipendula hexapetala</i> Gilib.	2	3	2	3	5	-	4	4	3	
<i>Fragaria viridis</i> Duchense	2	3	3	3	4	-	3	5	4	
<i>Gagea pratensis</i> (Pers.) Dumort.	2	4	4	3	5	-	4	5	4	
<i>Galega officinalis</i> L.	4	3	4	3	4	-	3	5	3	
<i>Galium aparine</i> L.	3	3	5	3	4	-	3	4	3	
<i>G. pedemontanum</i> All.	2	2	2	3	3	-	3	5	3	
<i>G. tricorne</i> Stokes	2	3	2	3	5	-	3	4	3	
<i>G. verum</i> L.	2	4	2	3	5	-	4	4	4	
<i>Geranium columbinum</i> L.	3	3	3	3	3	-	4	4	3	
<i>G. molle</i> L.	2	3	4	3	3	-	4	4	3	
<i>G. sanguineum</i> L.	2	4	2	3	3	-	3	4	4	
<i>Glechoma hederacea</i> L.	3	3	3	3	4	-	3	4	3	
<i>Glyceria fluitans</i> (L.) R. Br.	5	4	3	4	5	-	4	3	3	
<i>G. maxima</i> (Hartm.) Holombg.	5	4	5	3	5	-	4	4	3	
<i>Glycyrrhiza echinata</i> L.	3	3	2	3	3	-	4	4	3	
<i>Gratiola officinalis</i> L.	5	3	3	4	5	+	3	4	4	
<i>G. officinalis</i> L. f. <i>inundata</i> (Kit in Schultes) Soó										
<i>Gypsophila muralis</i> L.	3	2	1	2	3	-	4	4	4	
<i>G. muralis</i> L. var. <i>muralis</i> f. <i>capillaris</i> Fick. et Schube										
<i>Heleocharis palustris</i> (L.) R.Br.	5	4	2	4	5	-	4	3	3	
<i>Heleochloa alopecuroides</i> (Pill. et Mitterp.) Host	4	3	3	3	3	-	4	5	3	
<i>H. schoenoides</i> (L.) Host	4	4	3	3	4	+	4	5	3	
<i>Helminthia echioides</i> (L.) Gärtn.	3	3	4	4	4	-	4	5	2	
<i>Herniaria incana</i> Lam.	1	3	2	2	3	-	4	3	4	

Supplement – cont.

Taxon	Ecological index								
	F	R	N	H	D	S	L	T	K
<i>Hibiscus trionum</i> L.	3	3	3	3	4	-	4	5	4
<i>Hieracium pilosella</i> L.	2	3	2	3	4	-	4	3	4
<i>Holosteum umbellatum</i> L.	1	3	3	3	3	-	4	5	3
<i>Hordeum maritimum</i> subsp. <i>gussoneanum</i> (Parl.) A. et G.	2	4	3	3	5	+	5	4	3
<i>H. murinum</i> L.	2	3	4	2	3	-	4	4	4
<i>Hypericum perforatum</i> L.	2	3	3	3	5	-	3	4	3
<i>Inula britannica</i> L.	4	4	3	3	4	+	3	5	3
<i>Juncus articulatus</i> L.	4	3	2	3	5	-	4	3	3
<i>J. atratus</i> Krock.	4	2	3	4	5	-	4	4	2
<i>J. bufonius</i> L.	4	2	3	3	5	-	4	3	3
<i>J. compressus</i> Jacq.	4	3	3	3	5	+	4	3	3
<i>J. compressus</i> Jacq. var. <i>coartactus</i> E. Meyer ex Busch.									
<i>J. compressus</i> Jacq. var. <i>compressus</i> f. <i>porphyrocarpus</i> J. Murr.									
<i>J. conglomeratus</i> L.	4	2	3	4	5	-	4	3	2
<i>J. gerardi</i> Lois.	4	3	3	3	4	+	5	3	4
<i>Kickxia elatine</i> (L.) Dum.	2	3	3	3	4	-	4	5	2
<i>Knautia arvensis</i> (L.) Coult.	2	3	3	3	4	-	4	4	3
<i>Kochia laniflora</i> (Gmel.) Borb.	2	4	1	3	3	-	4	4	5
<i>K. prostrata</i> (L.) Schrad.	1	3	4	2	3	+	5	5	4
<i>K. prostrata</i> (L.) Schrad. l. <i>rubens</i> Lag.									
<i>Koeleria gracilis</i> Pers.	1	3	2	3	3	-	4	3	3
<i>K. gracilis</i> f. <i>violacea</i> R. et Sch.									
<i>Lactuca saligna</i> L.	1	3	4	3	4	+	4	5	4
<i>L. serriola</i> L.	2	3	3	2	3	-	4	5	4
<i>Lamium amplexicaule</i> L.	2	3	4	4	4	-	4	3	3
<i>L. purpureum</i> L.	3	4	4	3	4	-	4	3	3
<i>Lathyrus aphaca</i> L.	2	3	3	3	4	-	3	4	4
<i>L. aphaca</i> L. var. <i>aphaca</i> f. <i>laetus</i> Posp.									
<i>L. hirsutus</i> L.	2	4	3	3	3	-	3	4	4
<i>L. tuberosus</i> L.	2	4	3	3	5	-	4	4	4
<i>Leonurus cardiaca</i> L.	2	4	5	3	4	-	4	4	4
<i>L. marrubiastrum</i> L.	3	4	4	3	3	-	3	5	2
<i>Lepidium campestre</i> (L.) R.Br.	2	4	4	2	4	-	4	3	3
<i>L. draba</i> L.	2	3	4	3	4	-	3	4	4
<i>L. perfoliatum</i> L.	3	4	3	3	5	-	4	5	4
<i>L. ruderale</i> L.	2	3	4	2	3	-	4	4	4
<i>L. ruderale</i> L. f. <i>humile</i> Roch.									
<i>Linaria genistifolia</i> (L.) Mill.	2	4	2	3	3	-	4	4	4
<i>L. vulgaris</i> Mill.	2	3	3	3	4	-	4	4	3
<i>Lithospermum officinale</i> L.	3	4	3	3	4	-	3	4	3
<i>Lolium perenne</i> L.	3	3	4	3	5	-	4	3	3
<i>Lotus corniculatus</i> L.	2	4	3	3	4	-	4	3	3
<i>Lotus corniculatus</i> L. var. <i>pilosus</i> (Jord.) Posp.									

Supplement – cont.

Taxon	Ecological index								
	F	R	N	H	D	S	L	T	K
<i>L. tenuis</i> Kit.	3	4	2	3	5	+	4	4	2
<i>Lycopus europaeus</i> L.	5	3	3	5	5	-	3	4	3
<i>L. europaeus</i> L. f. <i>turfosus</i> Beck									
<i>L. exaltatus</i> L.	5	4	4	4	5	-	3	5	4
<i>Lysimachia nummularia</i> L.	4	3	4	3	5	-	2	4	2
<i>L. vulgaris</i> L.	4	3	3	4	5	-	3	4	3
<i>Lythrum hyssopifolia</i> L.	4	2	3	3	5	+	4	5	3
<i>L. salicaria</i> L.	4	3	3	4	5	-	3	4	3
<i>L. virgatum</i> L.	4	3	4	4	5	-	3	4	4
<i>Marrubium peregrinum</i> L.	2	4	2	2	5	-	4	4	4
<i>M. vulgare</i> L.	2	4	5	3	4	-	4	5	4
<i>Matricaria chamomilla</i> L.	3	3	3	3	4	+	4	4	3
<i>M. chamomilla</i> L. f. <i>salina</i> (Schur) Jáv.									
<i>M. inodora</i> L.	3	3	4	4	4	+	4	3	3
<i>M. suaveolens</i> (Pursh) Buch.	3	3	5	3	5	-	4	4	3
<i>Medicago falcata</i> L.	2	4	2	3	3	-	4	4	4
<i>M. lupulina</i> L.	2	4	3	3	4	-	3	4	3
<i>M. lupulina</i> L. f. <i>canescens</i> (Menth.) Soó									
<i>Melilotus officinalis</i> (L.) Pallas	2	4	3	3	3	-	4	4	3
<i>Mentha aquatica</i> L.	5	3	3	4	5	-	3	4	3
<i>Mentha aquatica</i> L. f. <i>erromera</i> Top.									
<i>M. longifolia</i> (L.) Huds.	4	4	4	3	5	-	3	3	3
<i>M. pulegium</i> L.	4	3	4	4	5	+	3	5	2
<i>M. pulegium</i> L. f. <i>lactea</i> Borb.									
<i>Muscari racemosum</i> (L.) Mill.	2	4	3	3	4	-	4	4	3
<i>Myosotis caespitose</i> (Schultz.)	5	2	3	3	4	-	4	4	3
<i>M. collina</i> Hoffm.	1	3	2	2	3	-	4	4	4
<i>M. palustris</i> (L.) Nath.	4	3	4	4	5	-	3	3	2
<i>Myosurus minimus</i> L.	4	2	2	3	5	-	4	4	2
<i>Nigella arvensis</i> L.	2	4	3	3	3	-	4	4	2
<i>Odontites rubra</i> Gilib.	3	3	4	3	5	+	3	4	3
<i>Oenanthe aquatica</i> (L.) Poir. in Lam.	5	4	3	3	5	-	4	4	4
<i>O. silaifolia</i> M.B.	5	4	3	3	5	+	4	4	3
<i>Oenothera biennis</i> L.	2	3	3	3	4	-	4	4	3
<i>O. strigosa</i> (Rydb.) Mach. et Bush	3	3	3	3	4	-	4	4	3
<i>Ononis spinosa</i> L.	2	3	2	3	5	-	4	4	3
<i>Onopordon acanthium</i> L.	2	4	5	3	4	-	4	4	4
<i>Ornithogalum gussonei</i> Ten.	2	4	3	4	4	-	4	5	4
<i>O. nutans</i> L.	3	3	3	3	4	-	4	5	2
<i>O. umbellatum</i> L.	3	4	3	3	4	-	4	4	2
<i>Orobanche caryophyllacea</i> Sm.	2	4	2	3	3	-	4	4	4
<i>Panicum crus-galli</i> L.	3	3	5	3	4	-	3	4	3
<i>Papaver dubium</i> L.	2	2	3	3	4	-	3	4	4

Supplement – cont.

Taxon	Ecological index								
	F	R	N	H	D	S	L	T	K
<i>P. rhoeas</i> L.	2	4	3	3	4	-	3	4	3
<i>Parentucellia latifolia</i> (L.) Car.	3	3	2	3	3	+	4	3	2
<i>P. latifolia</i> (L.) Car. f. <i>albiflora</i> (Raul.) Dunjić									
<i>Pastinaca sativa</i> L.	2	4	3	3	4	-	4	4	3
<i>Peplis portula</i> L.	4	2	3	3	5	-	4	4	2
<i>Petrorhagia prolifera</i> (L.) P. W. Ball.	1	2	2	3	3	-	4	5	3
<i>Peucedanum officinale</i> L.	2	4	2	3	5	-	3	5	3
<i>Pholiurus pannonicus</i> (Host) Trin.	3	4	2	3	5	+	4	4	4
<i>Phragmites communis</i> Trin.	5	3	3	3	4	+	3	3	3
<i>Picris hieracioides</i> L.	2	4	4	2	4	-	4	4	3
<i>Pimpinella saxifraga</i> L.	2	3	2	3	4	-	4	3	4
<i>Plantago altissima</i> L.	4	3	2	3	4	-	4	4	3
<i>P. lanceolata</i> L.	2	3	3	3	4	-	3	3	3
<i>P. lanceolata</i> L. var. <i>lanceolata</i>									
<i>P. lanceolata</i> L. var. <i>sphaerostachya</i> M.etK.									
<i>P. major</i> L.	3	3	4	3	5	+	4	3	3
<i>P. maritima</i> L.	2	4	2	2	4	+	3	3	5
<i>P. media</i> L.	2	4	2	3	4	-	4	3	3
<i>P. schwarzenbergiana</i> Schur.	3	3	3	2	5	+	4	4	4
<i>P. tenuiflora</i> W. et K.	3	4	1	3	5	+	4	3	4
<i>Poa annua</i> L.	3	3	4	3	4	-	4	3	3
<i>P. bulbosa</i> L.	1	3	3	2	3	-	5	4	4
<i>P. bulbosa</i> L. f. <i>vivipara</i> Koel.									
<i>P. pratensis</i> L.	3	3	3	4	4	-	4	3	3
<i>P. trivialis</i> L.	3	3	4	3	4	-	3	3	3
<i>Podospermum canum</i> C.A. Mey.	2	4	2	2	4	+	4	4	3
<i>P. canum</i> C.A. Mey. var. <i>integrifolia</i> (Schur) Soó									
<i>P. canum</i> C.A. Mey. var. <i>tenuissima</i> Borb.									
<i>Polygonum arenarium</i> W. et K.	1	3	1	2	2	-	4	4	4
<i>P. aviculare</i> L.	3	3	4	3	5	-	4	3	3
<i>P. aviculare</i> L. subsp. <i>aequale</i> (Lindm.) A. et G.	3	3	3	3	4	-	4	3	3
<i>P. lapathifolium</i> L.	3	3	4	3	3	-	5	3	3
<i>P. lapathifolium</i> L. var. <i>tomentosum</i> (Schrk.) Bay.									
<i>P. mite</i> Schrank	4	3	4	3	4	-	3	4	2
<i>P. patulum</i> M.B.	2	4	2	2	4	+	4	4	4
<i>P. persicaria</i> L.	3	3	4	3	3	-	4	3	3
<i>Portulaca oleracea</i> L.	3	3	4	3	4	-	4	4	3
<i>Potentilla anserina</i> L.	3	3	4	3	5	+	4	3	3
<i>P. argentea</i> L.	1	2	2	2	3	-	4	3	4
<i>P. argentea</i> L. var. <i>tenuiloba</i> (Jord.) M. Gusuleac.									
<i>P. leucopolitana</i> Müll.	1	3	2	2	3	-	4	5	4
<i>P. reptans</i> L.	3	3	4	3	5	-	4	3	3
<i>P. supina</i> L.	3	4	4	3	5	-	4	5	3
<i>Prunella vulgaris</i> L.	3	3	3	3	4	-	4	3	3

Supplement – cont.

Taxon	Ecological index								
	F	R	N	H	D	S	L	T	K
<i>Prunus spinosa</i> L.	2	4	3	3	3	-	4	4	3
<i>P. spinosa</i> L. f. <i>acuminata</i> (Clab.) Dom.									
<i>Puccinellia limosa</i> (Schur) Holmb.	3	4	2	2	4	+	4	4	4
<i>Pulicaria vulgaris</i> Gärtn.	4	2	5	3	5	+	4	5	3
<i>Pycnopus flavescens</i> (L.) Rchb	4	3	3	3	3	-	4	4	2
<i>Ranunculus acer</i> L.	3	3	3	3	4	-	3	3	4
<i>R. aquatilis</i> L.	5	3	4	4	5	-	3	4	2
<i>R. arvensis</i> L.	2	4	3	3	4	-	3	4	2
<i>R. lateriflorus</i> DC.	4	3	1	3	5	+	4	4	4
<i>R. paucistamineus</i> Tsch.	5	3	4	3	5	-	4	3	3
<i>R. pedatus</i> W. et K.	5	2	2	3	5	-	3	4	2
<i>R. repens</i> L.	4	3	4	3	5	-	3	3	3
<i>R. sardous</i> Cr.	4	3	3	3	5	+	4	4	2
<i>R. sceleratus</i> L.	4	4	5	4	5	+	4	4	3
<i>Rapistrum perenne</i> (L.) All.	2	3	4	3	4	-	3	5	4
<i>Rhinanthus borbasii</i> (Dörf.) Soó subsp. <i>borbasii</i>	2	4	3	3	3	+	3	3	5
<i>Roripa austriaca</i> (Cr.) Bess.	4	3	4	3	3	-	3	4	4
<i>R. kernerii</i> Menyh.	4	4	4	3	5	+	4	4	3
<i>Rubus caesius</i> L.	4	3	4	3	4	-	2	4	3
<i>Rumex crispus</i> L.	3	3	4	2	4	+	4	3	3
<i>R. obtusifolius</i> L.	3	3	4	4	4	-	4	3	3
<i>R. palustris</i> Sm.	4	4	4	3	5	-	4	5	3
<i>R. patientia</i> L.	3	3	4	3	4	-	4	5	4
<i>R. stenophyllus</i> Ledeb.	3	4	2	3	3	+	4	3	3
<i>Salsola ruthenica</i> Iljin	3	3	4	2	3	+	5	5	4
<i>S. soda</i> L.	3	4	4	3	4	+	4	4	4
<i>Salvia nemorosa</i> L.	2	3	4	3	4	-	4	5	4
<i>S. verticillata</i> L.	2	4	4	3	3	-	4	4	4
<i>Saponaria officinalis</i> L.	2	4	4	3	3	-	3	4	3
<i>Scabiosa ochroleuca</i> L.	1	4	2	3	3	-	4	5	5
<i>Schoenoplectus lacuster</i> (L.) Palla	5	3	3	4	4	-	5	4	3
<i>S. lacuster</i> (L.) Palla f. <i>conglomeratus</i> (Junge) Soó									
<i>Scilla autumnalis</i> L.	1	4	2	3	3	-	4	5	2
<i>Scleranthus annus</i> L.	3	1	2	3	3	-	4	4	4
<i>S. annus</i> L. f. <i>minimus</i> Schur.									
<i>Sclerochloa dura</i> (L.) Beauv.	1	3	3	2	5	-	5	5	4
<i>Scleropoa rigida</i> (L.) Griseb.	1	4	3	2	3	-	5	5	2
<i>Scutellaria hastifolia</i> L.	5	3	4	5	5	-	3	5	3
<i>Sedum caespitosum</i> (Cav.) DC.	1	4	1	2	4	+	5	4	3
<i>Senecio jacobea</i> L.	3	3	3	3	4	-	4	4	2
<i>S. vernalis</i> W. et K.	2	3	3	2	3	+	4	4	4
<i>Setaria glauca</i> (L.) P.B.	2	3	4	2	3	-	4	4	3
<i>S. viridis</i> (L.) P.B.	2	3	4	2	4	-	4	4	3



Supplement – cont.

Taxon	Ecological index									
	F	R	N	H	D	S	L	T	K	
<i>Silene alba</i> (Mill.) Krause	2	3	4	3	3	-	4	4	4	
<i>S. multiflora</i> (Ehrh.) Pers.	3	4	4	3	3	+	4	4	4	
<i>S. viscosa</i> (L.) Pers.	2	4	3	3	3	+	4	4	4	
<i>S. vulgaris</i> (Moench) Garcke	2	3	2	3	3	-	3	3	3	
<i>Sinapis arvensis</i> L.	3	4	4	3	4	-	4	4	3	
<i>Sisymbrium orientale</i> L.	2	3	4	3	4	-	4	5	2	
<i>Sonchus arvensis</i> L.	3	3	4	4	4	+	3	4	3	
<i>S. arvensis</i> L. var. <i>uliginosus</i> (M.B.) Grec.										
<i>S. asper</i> (L.) Hill.	3	3	4	3	4	-	4	4	3	
<i>Spergularia media</i> (L.) Presl.	3	3	3	3	3	+	4	4	3	
<i>S. salina</i> J. et C. Presl.	3	3	3	3	4	+	4	4	3	
<i>Stachys germanica</i> L.	1	4	3	3	4	-	4	5	3	
<i>Statice gmelini</i> Willd. subsp. <i>hungaricum</i> (Klokov) Soó	2	3	1	2	4	+	4	4	4	
<i>S. gmelini</i> Willd. subsp. <i>hungaricum</i> (Klokov) Soó f. <i>acuminatum</i> (Schur) Soó										
<i>S. gmelini</i> Willd. subsp. <i>hungaricum</i> (Klokov) Soó f. <i>obtusum</i> (Schur) Soó										
<i>S. gmelini</i> Willd. subsp. <i>hungaricum</i> (Klokov) Soó f. <i>hungaricum</i> (Schur) Soó										
<i>Symphytum officinale</i> L.	3	3	4	4	4	-	3	4	3	
<i>Taraxacum officinale</i> Weber	3	3	4	3	4	+	4	3	3	
<i>Taraxacum serotinum</i> /W.et.K./Poir. subsp. <i>bessarabicum</i> /Horn./ Hand.-Mazz.	3	4	1	3	4	+	4	4	4	
<i>Teucrium scordium</i> L.	5	4	3	5	5	+	4	5	3	
<i>Thalictrum lucidum</i> L.	4	4	3	4	5	-	3	4	4	
<i>Thesium linophyllum</i> L.	1	4	2	3	3	-	3	4	4	
<i>Thlaspi arvense</i> L.	3	3	4	4	4	-	3	3	3	
<i>Th. perfoliatum</i> L.	2	4	3	3	4	-	4	4	3	
<i>Thymus glabrescens</i> Willd.	1	4	2	2	3	-	4	5	5	
<i>Th. marschallianus</i> Willd.	1	3	1	3	2	-	4	4	4	
<i>Th. serpyllum</i> L.	2	3	1	2	3	-	4	2	4	
<i>Torilis arvensis</i> (Huds.) Link.	2	4	4	2	4	-	4	5	3	
<i>Trifolium angulatum</i> W.et K.	3	4	2	3	4	+	4	4	3	
<i>T. arvense</i> L.	1	2	1	2	3	-	4	4	4	
<i>T. campestre</i> Schreb.	2	3	2	3	4	-	4	4	3	
<i>T. campestre</i> Schreb. var. <i>minus</i> (Koch) Gremli										
<i>T. filiforme</i> L.	2	3	2	3	3	-	4	5	2	
<i>T. fragiferum</i> L.	3	4	2	3	5	+	4	4	3	
<i>T. hybridum</i> L.	3	4	4	3	4	-	4	3	3	
<i>T. ornithopodioides</i> (L.) Sm.	1	3	1	2	3	+	4	4	2	
<i>T. parviflorum</i> Ehrh.	1	3	2	2	4	+	4	4	3	
<i>T. pratense</i> L.	3	3	3	3	4	-	3	3	3	
<i>T. repens</i> L.	3	3	4	3	5	+	4	3	3	
<i>T. repens</i> L. f. <i>microphyllum</i> Larg.-Fossat										
<i>T. scabrum</i> L.	1	4	1	3	3	-	4	5	3	

Supplement – cont.

Taxon	Ecological index									
	F	R	N	H	D	S	L	T	K	
<i>T. striatum</i> L.	2	2	2	2	3	+	4	5	2	
<i>T. strictum</i> (L.) Jusl.	1	3	1	3	3	+	4	4	2	
<i>T. subterraneum</i> L.	1	2	2	3	3	-	4	5	2	
<i>T. vesiculosum</i> Savi.	1	3	1	3	3	+	4	4	3	
<i>Turgenia latifolia</i> (L.) Hoffm.	2	4	4	3	4	-	4	4	3	
<i>Typha angustifolia</i> L.	5	4	3	3	5	+	4	4	3	
<i>T. latifolia</i> L.	5	3	4	3	5	-	4	4	3	
<i>T. laxmannii</i> Lepech.	5	3	3	3	5	-	4	4	3	
<i>Typhoides arundinacea</i> (L.) Mnch.	5	3	4	3	4	-	3	4	3	
<i>Valerianella locusta</i> (L.) Betcke	3	3	4	3	4	-	4	4	2	
<i>Ventenata dubia</i> (Leers.) F. Schultz	1	3	2	2	3	-	3	5	3	
<i>Verbascum blattaria</i> L.	2	4	4	3	3	+	4	5	3	
<i>V. phoeniceum</i> L.	1	4	3	3	3	-	4	5	3	
<i>Verbena officinalis</i> L.	3	3	4	3	5	-	4	4	3	
<i>Veronica anagallis-aquatica</i> L.	5	3	4	4	5	-	4	4	3	
<i>V. anagallis-aquatica</i> L. f. <i>angustifolia</i> Krösche										
<i>V. anagalloides</i> Guss.	5	3	4	4	5	-	4	5	2	
<i>V. arvensis</i> L.	3	3	4	3	4	-	3	4	4	
<i>V. prostrata</i> L.	2	3	3	3	3	-	4	4	5	
<i>Vicia angustifolia</i> L.	2	3	3	3	4	-	4	5	4	
<i>V. hirsuta</i> (L.) S.F. Gray.	2	3	2	3	4	-	4	3	3	
<i>V. striata</i> M. Bieberst.	2	4	3	3	3	-	3	5	3	
<i>Viola arvensis</i> Murr.	3	3	3	3	3	-	3	3	3	
<i>Vulpia myuros</i> (L.) Gmel.	1	2	3	2	3	-	4	5	3	
<i>Xanthium italicum</i> Moretti	3	3	5	3	2	+	4	5	3	
<i>X. spinosum</i> L.	1	3	4	3	3	-	4	5	4	

Received: February 18<sup>th</sup> 2008  
 Revised: December 30<sup>th</sup> 2008  
 Accepted: December 30<sup>th</sup> 2008

## Pollen size and viability in hybrid swarm populations of *Pinus mugo* Turra and *Pinus sylvestris* L.

ANDREJ KORMUŤÁK<sup>1</sup>, BOŽENA VOOKOVÁ<sup>1</sup>, VLADIMÍR ČAMEK<sup>2</sup>, PETER BOLEČEK<sup>2</sup> & DUŠAN GÖMÖRY<sup>3</sup>

<sup>1</sup>Institute of Plant Genetics and Biotechnology, Slovak Academy of Sciences, Akademická A, P.O.Box 39A, SK-950 07 Nitra, Slovak Republic; e-mail: nrgrkorm@savba.sk

<sup>2</sup>Constantine Philosopher University, Faculty of Natural Sciences, A. Hlinku 1, SK-949 74 Nitra, Slovak Republic; vcamek@ukf.sk

<sup>3</sup>Technical University in Zvolen, Faculty of Forestry, T. G. Masaryka 24, SK-960 53 Zvolen, Slovak Republic; gomory@vsld.tuzvo.sk

---

Kormuťák A., Vooková B., Čamek V., Boleček P. & Gömöry D. (2008): Pollen size and viability in hybrid swarm populations of *Pinus mugo* Turra and *Pinus sylvestris* L. – Thaiszia – J. Bot. 18: 93-100. – ISSN 1210-0420.

Abstract: Pollen size and viability in two hybrid swarm populations of *Pinus mugo* and *P. sylvestris* were evaluated using microscopic approach and in vitro germination test. Pollen size of the hybrid swarms was shown to be reduced in comparison with the pollen size of the adjoining control populations of *P. mugo* and *P. sylvestris*. The differences in pollen germination percentage of hybrid swarms at Habovka and Suchá Hora and control population of *P. mugo* at Roháče have not reached the level of statistical significance. Statistically significant were only differences in pollen tube length. Hybrid swarms exhibited reduced pollen tubes of 15 % and 31 % in comparison with control populations. This indicates reduced vitality of pollen in hybrid swarms. The conclusion has been drawn postulating partial rather than complete sterility of the hybrid swarms.

Keywords: *Pinus mugo*, *P. sylvestris*, hybrid swarms, pollen, size, viability.

---

## Introduction

The two subgenera of the genus *Pinus* seem to be genetically differentiated not only with respect to species crossability but also with regard to fertility of their interspecific hybrids. The soft pines (subgenus *Haploxylon* and/or *Strobus*) are very prone to hybridize mutually with embryo inviability acting as the principal reproductive barrier between species (KRIEBEL 1972). On the contrary, the hard pines (subgenus *Diploxylon* and/or *Pinus*) are much more differentiated genetically involving gametophytic incompatibility as a powerful mechanism of reproductive isolation between species (BUCHHOLZ 1944, MCWILLIAM 1959). The consequence of a higher hybridological affinity among soft pines is a higher degree of fertility of their interspecific hybrids than in the hard pines. SAX (1960) has reported of 11 % of pollen sterility in *P. griffithii* × *P. strobus* and of 30 % of sterile pollen in *P. parviflora* × *P. strobus* hybrids of the soft pines as compared with the 40-50 % share of sterile pollen in the hard pine hybrids of *P. contorta* × *P. banksiana*. We have also found considerable reduction of pollen viability in the two hybrid swarm populations of *P. mugo* and *P. sylvestris* in northern Slovakia reaching the level of 40-41 % of sterile pollen (KORMUŤÁK et al. 2007).

In continuation with these experiments the study on pollen viability has been undertaken involving both pollen size and pollen germinability parameters in some other populations of the hybrid swarms and of *P. mugo* species in the region. The objective of the study was to validate the conclusions about reduced pollen viability in the hybrid swarms on broader experimental and geographical bases. This aspect of *P. mugo* and *P. sylvestris* hybrid swarm populations has until now been neglected, the attention being preferentially oriented towards morphometric and anatomic traits of the needles (BUSINSKY 1998, BORATYŇSKA et al. 2003).

## Materials and methods

### Material

The study on pollen viability has included the hybrid swarm populations of Scots pine (*Pinus sylvestris* L.) and Swiss pine (*P. mugo* Turra) at Habovka and Suchá Hora in the western part of the High Tatras. The population of *P. mugo* at Roháče adjoining the hybrid swarm at Habovka was used as a control. The location of individual populations is given in Tab. 1. The pollen samples of 30 trees of *P. mugo* at Roháče, 15 trees of putative hybrids at Habovka and 27 trees of putative hybrids at Suchá Hora were collected in spring 2007. Mature microstrobili of individual trees were harvested shortly before shedding of pollen and then transferred to the laboratory. Pollen was extracted from desiccated microstrobili by sieving. The dry pollen was stored in a desiccator over silica gel at 4°C for the period of 3 weeks and then used in germination test.

**Tab. 1. Species and populations used in the experiment**

Species/Hybrids	Locality	Altitude	Latitude	Longitude	Collect. year
<i>P. sylvestris</i>	O.Biely Potok	644 m	49°17'12"	19°33'18"	2006
<i>P. mugo</i>	Štrbské Pleso	1355 m	49°07'10"	20°03'41"	2006
Hybrid swarm	Habovka	815 m	49°16'25"	19°37'14"	2006
<i>P.mugo</i>	Roháče	1562-1720 m	49°12'27"	19°44'30"	2007
Hybrid swarm	Habovka	815 m	49°16'25"	19°37'14"	2007
Hybrid swarm	Suchá Hora	765 m	49°23'20"	19°47'11"	2007

***In vitro* germination**

Pollen germination was tested at 25°C on medium consisting of 1.5 % agar and 10 % sucrose. Each sample was triplicated. After 48 h the number of germinating pollen was recorded from a sample of 100 pollen grains, whereas the pollen tube length in a sample of 30 pollen grains of each Petri dish.

**Pollen size measurement**


In study on pollen size variation a 1-year stored pollen was used following its acetolysis in a mixture of acetic acid and sulphuric acid (ERDTMAN 1943). Study involved the pollen samples which originated from 12 trees of each the hybrid swarm population at Habovka, *P.mugo* population at Štrbské Pleso in the central part of the High Tatras and *P. sylvestris* population at Oravský Biely Potok adjoining the hybrid swarm at Habovka (Tab. 1). The measurement of the pollen corpus breadth and height was done microscopically using 50 pollen grains of each tree (Fig. 1).

**Statistics**

The differences in pollen size and in pollen tube length and germination percentage between localities and trees were tested by nested ANOVA. Because the percentages were bimodally distributed, the germination data were transformed using the arcsin transformation ( $p' = \arcsin \sqrt{p}$ ). All calculations were done using the GLM procedure of SAS (SAS 1988).

**Results**

Comparative study on pollen size revealed substantial differences between *P. sylvestris* and *P. mugo* species. *P. mugo* pollen was shown to be larger than pollen grains of *P. sylvestris* (Tab. 2) but the hybrid swarm population at Habovka possesses the smallest pollen grains approaching in this respect *P. sylvestris* rather than *P. mugo*. No differences between *P. sylvestris* and *P. mugo* were found in the pollen corpus breadth/height ratio both species sharing the same ratios. Having little lowered ratio the hybrid swarm at Habovka has however deviated from the parental species in this parameter.



**Fig. 1. Acetolysed pollen grains of *P. mugo* (left) and schematic drawing of pollen grain with traits measured (right).**

It follows from Tab. 3 that differences between investigated populations were statistically significant with regard to the pollen corpus breadth (F 7.02<sup>\*\*</sup>). The same was true of the pollen corpus height parameter (F 5.07<sup>\*</sup>). On the contrary, the differences in the pollen corpus breadth/pollen corpus height ratio between individual populations have not reached the level of statistical significance (F 1.55). It is worth of mentioning in this connection that all the three pollen size parameters investigated differed significantly between individual trees of the respective populations indicating profound individual variation in pollen size (Tab. 3).

**Tab. 2. Pollen size in hybrid swarm at Habovka and in populations of parental species**

Species/Hybrid	N	Pollen corpus breadth (µm) Mean ± SD	Pollen corpus height (µm) Mean ± SD	Breadth/height ratio
<i>P. sylvestris</i>	600	43.58 ± 3.91	33.75 ± 3.84	1.30 ± 0.14
Hybrid swarm	600	42.42 ± 4.23	33.41 ± 4.02	1.28 ± 0.15
<i>P. mugo</i>	600	45.47 ± 4.83	35.21 ± 4.52	1.30 ± 0.14

**Tab. 3. Variance analysis of pollen corpus breadth in hybrid swarm at Habovka and in parental species**

Source	D. f.	Sum of squares	Mean square	F value	Variance component [%]
Populations	2	2844.0	375278.0	7.02 <sup>**</sup>	21.0
Individuals	33	6681.3	32499.2	13.12 <sup>***</sup>	33.8
Error	1764	27222.4	400.1		39.9


Pollen viability tests has been preceded by a study on sucrose concentration effect in cultivation media. Among five concentrations tested the 10 % content of sucrose was found to be optimal for in vitro germination of *P. mugo* pollen (Tab. 4).

**Tab. 4. Effect of sucrose concentration in cultivation medium on germination of *P. mugo* pollen (The pooled data from two trees)**

Sucrose conc. [%]	N	Germination [%]	N	Pollen tube length [µm]
		Mean ± SD		Mean ± SD
3	600	84.16 ± 1.60	180	81.51 ± 23.58
5	600	84.33 ± 2.87	180	87.71 ± 22.25
10	600	92.00 ± 1.09	180	99.51 ± 26.13
13	600	82.16 ± 2.31	180	76.92 ± 27.11
15	600	79.00 ± 9.38	180	71.08 ± 17.84

N – sample size referring to germination percentage (600) and to pollen tube length (180)

As far as pollen viability of hybrid swarm populations at Habovka and Suchá Hora is concerned both populations were shown to produce less viable pollen than control population of *P. mugo* at Roháče. The total amount of 62.24 % and 67.21 % of germinating pollen in hybrid swarms represents reduction of pollen viability of 5 % in the hybrid population at Suchá Hora and of 12 % at Habovka in comparison with the pollen germinability in *P. mugo* (Tab. 5). In spite of these differences the variance analysis confirmed statistical significance of the differences at the individual level only (Tab. 6). The reduction of pollen germination potential in hybrid swarms has been paralleled by slowed pollen tube growth as evidenced by the average pollen tube length of 67.79 µm in hybrid swarm population at Habovka and of 55.44 µm length at Suchá Hora. The corresponding value in *P. mugo* population averaged at 80.41 µm deviating profoundly from the hybrid swarms (Tab. 7).



**Fig. 2. Germinating pollen of hybrid swarm at Suchá Hora (A) and *P. mugo* at Roháče (B)**

**Tab. 5. Pollen viability parameters in hybrid swarms and in *P. mugo***

Species/Hybrids	Locality	Germination [%]		Pollen tube length (µm)	
		N	Mean ± SD	N	Mean ± SD
<i>P. mugo</i>	Roháče	88	70.46 ± 17.96	2640	80.41 ± 27.78
Hybrid swarm	Habovka	45	62.24 ± 23.66	1350	67.79 ± 36.93
Hybrid swarm	Suchá Hora	84	67.21 ± 17.41	2520	55.44 ± 19.65

**Tab. 6. Variance analysis of pollen germinability in hybrid swarms and in *P. mugo* population**

Source	D.f.	Sum of squares	Mean square	F value
Populations	2	0.25999945	0.12999972	1.47
Individuals	13	1.15153096	0.08857930	34.37***
Repetitions	116	0.29898530	0.00257746	0.9534
Error	28	0.11460826	0.00409315	
Corr. total	216	9.87887122		

**Tab. 7. Variance analysis of pollen tube length in hybrid swarms and in *P. mugo* population**

Source	D.f.	Sum of squares	Mean square	F value	Variance component [%]
Populations	2	750556.0	375278.0	11.55***	21.0
Individuals	70	2274950.5	32499.2	81.22***	33.8
Error	6437	2575673.7	400.1		39.9
Corr. total	6509	5601180.2			

## Discussion

Among the aspects related to pollen viability in *P. sylvestris* and *P. mugo*, the studies aiming at optimizing cultivation media composition (CHIRA 1964), long-term storage of pollen (CHIRA 1971, OSTROLUCKÁ et al. 2003) and pollen viability variation at different stands in Slovakia (OSTROLUCKÁ et al. 1995) should be mentioned. Several studies have also appeared referring to pollen abortion and viability variation due to air pollution (FEDOTOV et al. 1983, FEDORKOV 1995, MIČIETA & MURIN 1998). Comparison of our experimental data with those published by other authors revealed some discrepancies which concern both characteristics studied, i. e. pollen size and pollen viability. CHIRA (1971) has for example reported of the non-acetolysed pollen corpus breadth ranging between 62.7  $\mu\text{m}$  and 85.8  $\mu\text{m}$  in *P. sylvestris* and between 62.7  $\mu\text{m}$  and 75.9  $\mu\text{m}$  in *P. mugo*. Taking into account relatively strong genetic control of this characteristic of the pollen, the observed discrepancy may be due to acetolysis treatment leading to destruction of the internal content of the pollen grains and to subsequent shrinking of the pollen corpus size. Obvious seem only the reasons of reduced pollen size in hybrid swarm populations which may be ascribed to the disturbed microsporogenesis in the swarms. It was shown in our previous study that increased frequency of meiotic disturbances occurring in hybrid swarm population at Habovka during anaphases I and II has resulted in a high frequency occurrence of aborted tetrads and aborted pollen grains leading to profound decline of pollen viability in 2005 and 2006 harvests (KORMUŤÁK et al. 2007). The same applies also for the 2007 harvest when the pollen germinability parameters of the hybrid swarms at Habovka and Suchá Hora have not reached the level of those ascertained in control population of *P. mugo* at Roháče. Especially it is true of the pollen tube length characteristics which may be taken as a convincing evidence of the lowered vitality of the hybrid swarms pollen.



Though statistically non-significant but optically conspicuous differences in germination percentage between hybrid swarms and control population of *P. mugo* may be ascribed to the detrimental effect of climatic conditions on microsporogenesis at high elevation of the locality Roháče (Tab. 1). A low amount of functional pollen under high mountain conditions is suggested to be very common (STERN & ROCHE 1974). In case of *P. mugo* population this amount has averaged at 70.46 % which is much lesser than pollen germinability observed at the localities Štrbské Pleso and Popradské Pleso which are located at lower elevations and where pollen germinability averaged during 2005-2006 at 82.83 % and 78.38 % respectively (KORMUŤÁK et al. 2007). It is worth of mentioning that CHIRA (1971) has reported of 60 % germinability of freshly collected pollen of *P. mugo* whose origin has not however been given. Using pollen samples from 7 localities of the High Tatras OSTROLUCKA & FLEISHER (1995) have illustrated both annual and site variations in germinability of *P. mugo* with germination percentage ranging between 48 % and 76 % and pollen tube length averaging at 82-123  $\mu\text{m}$ . Profound effect of climatic conditions on the course of microsporogenesis in forest trees and resulting annual variation in quality of mature pollen seem to be a reasonable explanation of the outlined differences in pollen viability of *P. mugo* at Štrbské Pleso and Popradské Pleso as well. According to degree of sterility STEBBINS (1950) divided interspecific hybrids of plants into two groups. The first group is represented by the hybrids capable of producing some viable pollen and seed set through selfing, intercrossing between  $F_1$  individuals or backcrossing to the parental species. The second group involves the hybrids which are completely sterile. Presented experimental data provided evidence of partial rather than complete sterility of the hybrid swarm population of *P. mugo* and *P. sylvestris*. This may be taken as an additional support for the conclusion of SAYLOR & SMITH (1966) about partially reduced reproductive capacity of the interspecific hybrids of hard pines.

## Acknowledgement

This work was supported by the Slovak Science and Technology Assistance Agency under contract No. APVT-51-004-004.

## References

- BORATYŇSKA K., BORATYŇSKI A. & LEWANDOWSKI A. (2003): Morphology of *Pinus uliginosa* (Pinaceae) needles from populations exposed to and isolated from the direct influence of *Pinus sylvestris*. Botanical Journal of the Linnean Society 142: 83-91.
- BUCHHOLZ J. T. (1944): The cause of sterility in cross-pollinations between certain species of pines. American Journal of Botany 31: 25-26.
- BUSINSKY R. (1998): *Pinus mugo* complex in former Czechoslovakia – taxonomy, distribution, hybrid populations and endangerment. Report of Czechoslovak Botanical Society, Prague 33: 29-52.
- CHIRA E. (1964): Effect of storage conditions and storage period on pollen vitality in some species of the genus *Pinus*. Sborník vysoké školy zemědělské v Brně. Řada C: Spisy fakulty lesnické 3: 149-159. [In Slovak]

- CHIRA E. (1971): Cytogenetic methods in forest tree breeding. *Príroda Bratislava*. [In Slovak]
- ERDTMAN G. (1943): An introduction to pollen analysis. Verdoorn, New Ser. Plant Sci. Books, 12, Waltham, Mass.
- FEDORKOV A. L. (1995): Microsporogenesis, male flowering and seed development of Scots pine trees under pollution. In: Korpilähti E., Salonen T. & Oja S. (eds.), *Caring for the Forest: Research in a Changing World: 102-103*, Poster Abstracts, IUFRO XX World Congress, 6-12 August 1995, Tampere, Finland.
- FEDOTOV I. S., KARABAN R. T., TIKHOMIROV F. A. & SISIGINA T. I. (1983): Evaluation of SO<sub>2</sub> effects on Scots pine stands. *Lesovedeniye* 6: 23-27. [In Russian]
- KORMUTAK A., BOHOVIČOVA J., VOOKOVA B. & GÖMÖRY D. (2007): Pollen viability in hybrid swarm populations of *Pinus mugo* Turra and *P. sylvestris* L. *Acta Biologica Cracoviensia Series Botanica* 49: 61-66.
- KRIEBEL H. B. (1972): Embryo development and hybridity barriers in the white Pines (Section *Strobus*). *Silvae Genetica* 21: 39-43.
- McWILLIAM J. R. (1959): Interspecific incompatibility in *Pinus*. *American Journal of Botany* 46: 425-433.
- MÍČIETA K. & MURÍN G. (1998): Three species of genus *Pinus* suitable as bioindicators of polluted environment. *Water, Air, and Soil Pollution* 104: 413-422.
- OSTROLUCKA M. G., ČIČOVA L. & BOLVANSKY M. (2003): Influence of temperature and storage on pollen viability in *Pinus mugo* Turra. *Folia Oecologica* 30: 157-162.
- OSTROLUCKA M. G. & FLEISHER P. (1995): Pollen vitality monitoring in the Tatra biosphere reserve. *Ekológia (Bratislava)* 14: 391-397.
- SAX K. (1960): Meiosis in interspecific pine hybrids. *Forest Science* 6: 135-138.
- SAYLOR L. C. & SMITH B. W. (1966): Meiotic irregularity in species and interspecific hybrids of *Pinus*. *American Journal of Botany* 53: 453-468.
- STEBBINS G. L. J. R. (1950): *Variation and evolution in plants*. Columbia University Press, New York, 643 pp.
- STERN K. & ROCHE L. (1974): *Genetics of forest ecosystems*. Springer-Verlag Berlin Heidelberg New York, 330 pp.

Received: January 18<sup>th</sup> 2008  
 Revised: December 30<sup>th</sup> 2008  
 Accepted: December 30<sup>th</sup> 2008

### Book review

---

**W. FREY & R. LÖSCH: Lehrbuch der Geobotanik. Pflanze und Vegetation in Raum und Zeit. 2. Auflage [Handbook of geobotany. Plants and vegetation in space and time].**

Springer/Spektrum Akademischer Verlag, Tiergartenstrasse 17, 69121 Heidelberg, Germany; 2004, XIV, 528 S., Geb., ISBN: 978-3-8274-1193-8, Price: \$59.95.  
<http://www.springer.com/spektrum+akademischer+verlag/>

The authors succeeded in creating the very actual and comprehensive textbook for students of geobotany. This second edition follows the last one in modern representation and it was fulfilled with some new, actual and, at the present time, necessary information for a geobotanist from parallel botanical disciplines. The textbook is well arranged and richly structured by its text, graphs, pictures, tables and schemes. The contents of teaching are divided into 10 chapters with other subchapters. After the first two introductory chapters, the third chapter Floristics and Areal presents floristic areals, cosmopolitanism and endemism, floristic geoelements, areal types, floristic structure of the Earth, biodiversity, species richness, phytogeographical areas and phytogeographical regions. The fourth chapter deals with geobotany itself, structure of communities, syntaxonomy, multivariable methods in phytocoenology, syntaxonomy and syndynamics. In the fifth chapter, the historical geobotany is discussed. The most space is devoted to plant ecology in the sixth chapter – from abiotic environmental conditions through autecology and ecological physiology, competitive relations to ecosystems and their modeling. In the seventh chapter, the ecological bases for populations and communities and reproductive ecology are presented. The eighth chapter deals with adaptations and strategies of living. The ninth chapter describes the individual vegetation zones and biomes in the Earth. The influence of human activities to biotopes, their conservation, risk to species and ecosystems are discussed in the tenth chapter. The textbook has a very rich list of literature, well arranged and informative graphs, schemes, tables and figures. At the beginning, there are explanatory notes to abbreviations, symbols and mathematical constants. At the end of the book, a comprehendious alphabetical dictionary of terms helps to use the textbook more quickly and easily.

I can recommend this textbook to university students, teachers, employers in nature protection, in state administration and self-administration dealing with environmental problems, foresters, farmers and to everyone which is interested in vegetation and its conservation. It should not be missing in any library of educative institutions and environmental organizations.

S. MOCHNACKÝ



## Professor VIRA V. PROTOPOPOVA – 75 Anniversary



In 2008 East European scientific society celebrates a jubilee of famous Ukrainian botanist Prof. VIRA V. PROTOPOPOVA, laureate of M.G. Kholodny Prize of National Academy of Sciences of Ukraine, Honorary member of Ukrainian Botanical Society and head scientist researcher at the M.G. Kholodny Institute of Botany, NAS of Ukraine. She is well known as specialist in floristics, plant geography and systematics. In Ukrainian botany the name of V.V. Protopopova is mainly associated with the scientific field of synanthropic flora, including such phenomenon as invasions of alien species. She is well known specialist in orchids and ferns systematic as well.

VIRA V. PROTOPOPOVA was born on 2 March 1933 in Kharkiv. Her father, Prof. Dr. VIKTOR P. PROTOPOPOV, was a famous scientist, teacher and organizer of medicine science, academician of Ukrainian Academy of Sciences, head of department of psychophysiology at the Institute of physiology Academy of Sciences of Ukraine, mother, RYMA V. SOBOLEVSKA, was a practical doctor.

V. PROTOPOPOVA begins her education in Kharkiv's school in 1941, and finished in Kiev's school in 1952. In this year, she entered the Taras Shevchenko State University and specialized in botany. Among her teachers were famous pedagogues and scientists, botanists, academicians DMYTRO K. ZEROV and OLEKSANDER V. TOPACHEVSKY, professors OLEXIY L. LYPA, associate professor PETRO M. BEREGOVYI, etc. During early students years VIRA PROTOPOPOVA starts to take part in scientific investigations, and prepared interesting works on systematic of *Lamiaceae* in Kaniv state reserve (Cherkasy Region), flora of Martyan Cape (the Crimea), and flora of Rzhyshev forest (Kyiv Region). In 1958 she finished the university with diploma on speciality „biology”.

In 1956-1958, VIRA V. PROTOPOPOVA worked at the Academician O.V. Fomin Botanical Garden of Taras Shevchenko State University. Since 1958 VIRA V. PROTOPOPOVA has worked in department of systematics and floristics of vascular plants of M.G. Kholodny Institute of Botany, Academy of Sciences of Ukraine: as assistant (1958-1961), post-graduate (1961-1964), junior scientific researcher (1964-1974), senior scientific researches (1974-1992), and head scientific researcher (1992- until now).

In 1966, under supervising Prof. MYKHAYLO I. KOTOV, VIRA V. PROTOPOPOVA prepared and defended her PhD degree work (candidate of biological sciences), with further publishing original monograph on this base („Alien plants of Forest-Steppe and Steppe of Ukraine ”, 1973). Later, in 1989, V. V. PROTOPOPOVA defended DrSc work (doctor of biological sciences), with publishing another monograph as well („Synanthropic flora of Ukraine and ways of its development”, 1991). In 1996, these monographs together with numerous papers under generalized title „Theoretical and practical aspects of natural differentiation and synanthropization of spontaneous flora of Ukraine” were awarded with M.G. Kholodny Prize of National Academy of Science of Ukraine.

We are to elucidate the main results obtained by Prof. V. V. PROTOPOPOVA now. She was pioneered in exploring of composition of synanthropic flora in Ukraine, defining its general and regional features, distribution of the main groups of synanthropic species. It is offered and proved a hypothesis of threshold of a species diversity of synanthropic floras with working out the index for estimation of ecological condition in different regions. Prof. V. V. PROTOPOPOVA also analyzed 150-year history of adventization of Ukrainian flora and estimated factors and revealed reasons of alien plants area dynamics in Ukraine. All this serves as a basis for putting into practice scientific forecast of synanthropization of regional floras depending on environmental changes. In addition, the general directions of synanthropization and adventization of Ukrainian flora are determined, and ecological and floristic classification of alien plants is developed. It is offered original hypothesis on origin of some local

endemics from archaeophytes, which opened new aspect in research of local endemism in anthropogenic transformed territory. She assumed that evolution of alien plants within the limits of secondary areas occurred through hybridization and allopolyploidisation with the subsequent differentiation of sympatric and marginal races.

Prof. VIRA V. PROTOPOPOVA prepared detail, critical and systematic investigations of many taxa of natural and alien fractions of Ukrainian flora, such as Lycopodiophyta, Equisetophyta, Polypodiophyta, *Orchidaceae*, *Amaranthaceae*, *Chenopodiaceae*, *Oxalidaceae*, some group of *Asteraceae* (*Bidens* L., *Galinsoga*, *Helianthus*, *Xanthium*, etc.). The results obtained were included in many Manuals and Floras, such as «Manual of plants of Ukraine» (1965), «Weeds of Ukraine» (1970), «Manual of plants of Ukrainian Carpathians» (1974), «Red Data Book of Ukraine» (1980, 1996), «Flora of European part of the USSR» (vol. 2 – 1976; vol. 4 – 1979; vol. 5 –1981), «Flora of Eastern Europe» (vol. 7 – 1994), «Manual of higher plants of Ukraine» (1987), «Chorology of Ukraine» (1987), «Ecoflora of Ukraine» (2002). and Prof. V.V. Protopopova is authors and co-authors more than 200 publications in many scientific journals, such as Ukrainian Botanical journal, Botanical journal, Biological sciences (Russia), Acta Botanica Slovaca, Thaiszia (Slovak Republic), Euphytica (USA), etc.

During her teacher's career Prof. VIRA V. PROTOPOPOVA supervised 3 dissertations of post-graduates: Drs. IRYNA A. TYMCHENKO ("Structure of population of species of tribe *Neottieae* Lindl. (*Orchidaceae*) Ukrainian flora and its tendencies of changes in condition of synanthropization", 1996), TETYANA V. VASYLYEVA-NEMERTSALOVA („Synanthropic flora of near port towns of North-Western of Black Sea area", 1996), and IRYNA V. KOVTUN („Flora of Kamyanets Cis-Dnister", 2004); last years she was a professor at Pereyaslav-Khmelnytsky state pedagogical university.

Special interest of VIRA V. PROTOPOPOVA connected with a popularization of botanical knowledge. Together with her colleagues TETYANA L. ANDRIYENKO, BORYS V. ZAVERUKHA, SVITLANA S. MOROZYUK, LIDIYA S. PANOVA she prepared many interesting, substantial, and poetic works: „Botanical album" (1979), „Protected plants of Ukraine" (1983), „Steppe plants" (1983), „Plants-travellers" (1989), „Herbaceous plants" (1986), „Spring plants" (1987), „Herbaceous plants of Ukraine" (2007), „Spring plants of Ukraine" (2007). These educational books got a wide public acknowledgement.

During a long time Prof. VIRA V. PROTOPOPOVA was a member of Botanical division of National Commission on Red Data Book of Ukraine, member of editorial board of Ukrainian Botanical journal, member of Qualification Council of DrSc degree at M.G. Kholodny Institute of Botany, NAS of Ukraine, Head of section of history of sciences of Ukrainian Botanical Society, and now she is a member of Qualification Council of DrSc degree at M.M. Gryshko National Botanical Garden of NAS of Ukraine.

Colleagues also known VIRA V. PROTOROVA as charming woman, with wide open soul and heart full of love to people.

Botanists, colleagues, and friends congratulate VIRA V. PROTOROVA with 75 jubilee. Long live, Dear Professor!

K. ROSTAŃSKI, A. TERPO, S. MOCHNACKY,  
M. SHEVERA, A. ILYNSKA, I. TYMCHENKO, A. YENA

### Selected publications of V.V. PROTOROVA\*

#### 1960

Котов М.І., Протопопова В.В. Знахідка кримської орхідеї –ремнепелюстника козлиного (*Hymanthoglossum caprinum* (M.V.) Spreng.) на Кавказі // Укр. ботан. журн. – 1960. – **17**, № 2.– С. 77–82 [Котов М.І., Протопопова В.В. 1960. New record of Crimean orchid – *Hymanthoglossum caprinum* (M.V.) Spreng. in Caucasus. Ukr. Bot. Journ. **17**, 2. Pp. 77–82].

#### 1963

Кузнецова Г.О., Протопопова В.В. Флора і рослинність острова Орлів на Чорному морі // Укр. ботан. журн. – 1963. – **20**, № 4.– С. 80–85 [KUZNETSOVA G.O., PROTOROVA V.V. 1963. Flora and vegetation of Black sea island Orliv. Ukr. Bot. Journ. **20**, 4. Pp. 80-85].

#### 1964

Протопопова В.В. Крейдяний бір на горах Артема та необхідність його охорони // Охороняйте рідну природу. III Зб. статей з питань охорони і раціонального використання природних ресурсів УРСР. – К.: Урожай, 1964. – С. 80–86. [PROTOROVA V.V. 1964. Cretaceous pineforest on Artem mountain and nessecity of its protection. To protect native nature. III Proceeding on questions of protection and use of natural resources of URSS. Pp. 80-86. Kyiv: Urozhay Press.].

Протопопова В.В. Поширення деяких нових та маловідомих адвентивних рослин по Лісостепу і Степу України за післявоєнний час // Питання експеримент. ботан. / Відпов. ред. К.М. Ситник. – 1964. – С. 127–132. [PROTOROVA V.V. 1964. Distribution of some new and little known alien plants in Forest-Steppe and Steppe of Ukraine for post-war time. Questions of experimental botany. Ed. K.M. SYTNIK. Pp. 80-86]

Протопопова В.В. Нові дані про систематичний склад роду нетреба (*Xanthium* L.) на Україні // Укр. ботан. журн. – 1964.– **21**, № 4.– С. 78–84 [PROTOROVA V.V. 1964. New data on systematic composition of the genus *Xanthium* L. in Ukraine. Ukr. Bot. Journ. **21**, 4. Pp. 78-84].

---

\* General list of Prof. V.V. PROTOROVA's publications is presented in A.P. ILYNSKA, I.A. TYMCHENKO and M.V. SHEVERA (2008) „VIRA VIKTORIVNA PROTOROVA”: [www.botany.kiev.ua/public.htm](http://www.botany.kiev.ua/public.htm)



#### 1965

- ПРОТОПОПОВА В.В. Адвентивні рослини Лісостепу та Степу України // Укр. ботан. журн. – 1965. – **22**, № 3.– С. 38–44 [PROTOROVA V.V. 1965. Alien plants of Forest-Steppe and Steppe of Ukraine. Ukr. Bot. Journ. **22**, 3. – Pp. 38-44].
- ПРОТОПОПОВА В.В. Родина Орхідних – *Orchidaceae* // Визначник рослин України. Вид. 2, виправ. і доповн. / Відпов. ред. Д.К. ЗЕРОВ. – К.: Урожай, 1965. – С. 177–186. [Protopyova V.V. 1965. *Orchidaceae*. Manual of plants of Ukraine. Ed. D.K. ZEROV. – Pp. 177–186. Kyiv: Urozhay Press.].

#### 1970

- ПРОТОПОПОВА В.В. *Commelinaceae, Cannabinaceae, Urticaceae, Amaranthaceae, Nyctaginaceae, Portulacaceae, Balsaminaceae, Thymelaeaceae, Onagraceae, Martyniaceae, Campanulaceae* Бур'яни України (Визначник-довідник) / Відпов. ред. О.Д. ВІСЮЛІНА. – К.: Наук. думка, 1970. – С. 65-66, 71-73, 112-118, 246-247, 254-256, 353-353, 372 [PROTOROVA V.V., 1965. *Commelinaceae, Cannabinaceae, Urticaceae, Amaranthaceae, Nyctaginaceae, Portulacaceae, Balsaminaceae, Thymelaeaceae, Onagraceae, Martyniaceae, Campanulaceae*. – Weeds of Ukraine. Ed. O.D. VISYULINA. – Pp. 65-66, 71-73, 112-118, 246-247, 254-256, 353-353, 372. Kyiv: Naukova dumka Press.].
- ПРОТОПОПОВА В.В., ДОБРОЧАЄВА Д.М., КУЗНЕЦОВА Г.О. Складноцвітні – *Compositae* // Бур'яни України (Визначник-довідник) / Відпов. ред. О.Д. ВІСЮЛІНА. – К.: Наук. думка, 1970. – С. 372–431 [PROTOROVA V.V., DOBRUCHAYEVA D.M., KUZNETSOVA G.O. *Asteraceae*. Weeds of Ukraine. Ed. O.D. VISYULINA. Pp. 372-431. Kyiv: Naukova dumka Press].

#### 1973

- ПРОТОПОПОВА В.В. Адвентивні рослини Лісостепу і Степу України / Відпов. ред. М.І. КОТОВ. – К.: Наук. думка, 1973. –192 с. [PROTOROVA V.V. 1973. Alien plants of Forest-Steppe and Steppe of Ukraine. Ed. M.I. KOTOV. 192 p. Kyiv: Naukova dumka Press].

#### 1974

- ПРОТОПОПОВА В.В. До видового складу квасениць (*Oxalis* L.) на Україні // Укр. ботан. журн. – 1974. – **31**, № 1.– С. 110–113 [PROTOROVA V.V. 1974. On the species composition of *Oxalis* L. genus in Ukraine. Ukr. Bot. Journ. **31**, 1. Pp. 110–113].
- ПРОТОПОПОВА В.В. Нові для України види роду *Diphysium* C. Presl. // Укр. ботан. журн. – 1974. – **31**, № 6.– С. 690–694 [PROTOROVA V.V. New for Ukraine species of the genus *Diphysium* C. Presl. Ukr. Bot. Journ. **31**, 1. Pp. 690–694].

#### 1976

- ПРОТОПОПОВА В.В. Семейство *Commelinaceae* – Комелиновые // Флора европейской части СССР / Отв. ред. АН.А. ФЕДОРОВ. – Л.: Наука, 1976. – Т. 2. – С. 219–220 [PROTOROVA V.V. 1976. Family *Commelinaceae*. Flora of European part of USSR. Ed. AN.A. FEDOROV. Vol. 2. Pp. 219-220. Leningrad: Nauka].

#### 1977

- ПРОТОПОПОВА В.В. Відділи Lycopodiophyta, Equisetophyta, Polypodiophyta, *Amaranthaceae, Saxifragaceae, Asteraceae* (*Ambrosia* L., *Galinsoga* L., *Xanthium* L., *Bidens* L., *Cyclachaena* Fresen.), *Orchidaceae*. – Визначник рослин Українських Карпат / Відпов. ред. В.І. ЧОПІК. – К.: Наук. думка, 1977. – С. 22–35,

101–104.147–152. 294–296. 382–390 [ПРОТОРОВА V.V. 1977. Lycopodiophyta, Equisetophyta, Polypodiophyta, *Amaranthaceae*, *Saxifragaceae*, *Asteraceae* (*Ambrosia* L., *Galinsoga* L. *Xanthium* L., *Bidens* L., *Cyclachaena* Fresen.), *Orchidaceae*. Manual of plants of Ukrainian Carpathians. Ed. V.I. СНОРУК. Pp. 22–35, 101–104.147–152. 294–296. 382–390. Kyiv: Naukova Dumka Press].

#### 1978

ПРОТОРОВА V.V. 1978. Adventive flora of the flat part of the Ukraine // Acta Botan. Slovaca. Ser. A (3). – Bratislava. – Pp. 71-74.

#### 1979

ПРОТОПОПОВА В.В. Семейство *Pontederiaceae* // Флора европейской части СССР / Под ред. А.А. ФЕДОРОВА. – Т. IV. – Л.: Наука, 1979. – С. 291–292. [ПРОТОРОВА V.V. 1979. Family *Pontederiaceae*. Flora of European part of USSR. Ed. А.А. ФЕДОРОВ. Vol. 4. Pp. 291-292. Leningrad: Nauka].

ПРОТОПОПОВА В.В., ТКАЧЕНКО В.С. Історія та прогноз поширення *Grindelia squarrosa* (Pursh) Dunal // Укр. ботан. журн. – 1979. – **36**, № 5. – С. 457–461 [ПРОТОРОВА V.V., ТКАЧЕНКО V.S. 1979. History and prognosis of distribution of *Grindelia squarrosa* (Pursh) Dunal. Ukr. Bot. Journ. **36**, 5. Pp. 457–461].

МОРОЗЮК С.С., ПРОТОПОПОВА В.В. Альбом з ботаніки (посібник для вчителів). – К.: Рад. шк., 1979. – 151 с. [МОРОЗЮК S.S., ПРОТОРОВА V.V. 1979. Botanical album (manual for teachers). 151 p. Kyiv: Radyanska shkola Press.]

#### 1980

ДУБИНА Д.В., ПРОТОПОПОВА В.В. Нові для флори СРСР види водяних папоротей з родини *Azollaceae* // Укр. ботан. журн. – 1980. – **37**, № 5. – С. 20–26 [ДУБИНА D.V., ПРОТОРОВА V.V. 1980. New species for USSR flora of water ferns from family *Azollaceae*. Ukr. Bot. Journ. **37**, 5. Pp. 20–26].

ПРОТОПОПОВА В.В. Родина Зозулинцеві // Червона книга Української РСР / Відпов. ред. К.М. Ситник. – К.: Наук. думка, 1980. – С. 252–253, 258–295, 318–331, 350–355, 362–363, 366–369, 372–397, 410–411, 414–415, 428–429, 436–437, 456–457, 476–477, 484–485, 490–491 [ПРОТОРОВА V.V. Family *Orchidaceae*. Red data Book of Ukraine. Ed. К.М. Sytnik. Pp. 252–253, 258–295, 318–331, 350–355, 362–363, 366–369, 372–397, 410–411, 414–415, 428–429, 436–437, 456–457, 476–477, 484–485, 490–491. Kyiv: Naukova dumka Press].

#### 1981

ДУБИНА Д.В., ПРОТОПОПОВА В.В. Знахідка *Pilularia globulifera* в УРСР // Укр. ботан. журн. – 1981. – **38**, № 6. – С. 56–59 [ДУБИНА D.V., ПРОТОРОВА V.V. 1981. Find of *Pilularia globulifera* in Ucr.RSS. Ukr. Bot. Journ. **38**, 6. Pp. 56–59].

ПРОТОПОПОВА В.В. Семейство *Martiniaceae* – Мартиниевые / Флора европейской части СССР / Под. ред. А.А. ФЕДОРОВА. – Т. V. – Л.: Наука, 1981. – С. 316–317 [ПРОТОРОВА V.V. 1981. Family *Martiniaceae*. Flora European part of USSR. Ed. А.А. ФЕДОРОВ. Vol. 5. Pp. 316–317. Leningrad: Nauka].

#### 1983

ДУБИНА Д.В., ПРОТОПОПОВА В.В. *Chenopodium pumilio* R. Br. – новий для флори СРСР вид // Укр. ботан. журн. – 1983. – **40**, № 4. – С. 47–51 [ДУБИНА D.V., ПРОТОРОВА V.V. 1983. *Chenopodium pumilio* R. Br. – new species on URSS flora. Ukr. Bot. Journ. **40**, 4. Pp. 47–51].

ДУБИНА Д.В., ПРОТОПОПОВА В.В. Эколого-биологические особенности *Azolla caroliniana* Willd. и *A. filiculoides* Lam. // Растит. ресурсы. – 1983. – **19**, вып. 4. – С. 500–506 [DUBINA D.V., PROTOROVA V.V. 1983. Ecological and biological peculiarities of *Azolla caroliniana* Willd and *A. filiculoides* Lam. Plant resources. **19**, 4. Pp. 500-506].

ПРОТОПОПОВА В.В. Ареалогическая характеристика орхидных Украины // Охрана и культивирование орхидей: Тез. докл. II Всесоюзн. совещ. (г. Киев, май 1983 г.) / Отв. ред. А.М. Гродзинский. – Киев: Наук. думка, 1983. – С. 112–113. [PROTOROVA V.V. 1983. Areal characteristic of Ukrainian orchids. Protection and cultivation of orchids: Abstracts of II All-Union conf. (Kiev, May, 1983). Ed. A.M. GRODZYNSKY. Pp. 112–113. Kyiv: Naukova dumka Press].

ЗАВЕРУХА Б.В., АНДРІЄНКО Т.Л., ПРОТОПОПОВА В.В. Охраняемые растения УССР. – Киев: Наук. думка, 1983. – 175 с. [ZAVERUKHA B.V., ANDRIYENKO T.L., PROTOROVA V.V. 1983. Protected plants of Ukraine. 173 p. Kyiv: Naukova dumka Press].

ПАНОВА Л.С., ПРОТОПОПОВА В.В. Степові рослини. – К.: Рад. шк., 1983. – 190 с. [PANOVA L.S., PROTOROVA V.V. 1983. Steppe plants. 190 p. Kyiv: Radyanska shkola Press].

#### 1984

ДУБИНА Д.В., ПРОТОПОПОВА В.В. *Euphorbia maculata* L. – новий для флори України вид // Укр. ботан. журн. – 1984. – **41**, № 3.– С. 33–36 [DUBINA D.V., PROTOROVA V.V. 1984. *Euphorbia maculata* L. – new for Ukraine species. Ukr. Bot. Journ. **41**, 3. Pp. 33–36].

ДУБИНА Д.В., ПРОТОПОПОВА В.В. Новий для флори європейської частини СРСР вид *Torulanium ferax* (Rich.) Urb. // Укр. ботан. журн. – 1984. – **41**, № 5.– С. 21–25 [DUBINA D.V., PROTOROVA V.V. 1984. New for European part of USSR species *Torulanium ferax* (Rich.) Urb. Ukr. Bot. Journ. **41**, 5. Pp. 21–25].

ПРОТОПОПОВА В.В. Особливості розподілу синантропних рослин за окремими ботаніко-географічними районами України // Укр. ботан. журн. – 1984. – **41**, № 1.– С. 46–49 [PROTOROVA V.V. 1984. Peculiarities of distribution of synanthropic plants in different botanical and geographical regions of Ukraine. Ukr. Bot. Journ. **41**, 1. Pp. 46–49].

#### 1985

ДУБИНА Д.В., ПРОТОПОПОВА В.В. Анализ адвентивной флоры заповедника „Дунайские плавни” // Биол. науки. – 1985. – № 10. – С. 68–73 [DUBINA D.V., PROTOROVA V.V. 1985. Analysis of alien flora of Dunaysky plavni reserve. Biol. Nauk. 10. Pp. 68–73].

#### 1986

ДУБИНА Д.В., ПРОТОПОПОВА В.В., ДУБОВИК О.М. Новий для УРСР рід *Brachyactis* Ledeb. // Укр. ботан. журн. – 1986. – **43**, № 2.– С. 51–54 [DUBINA D.V., PROTOROVA V.V., DUBONIK O.M. 1986. New for Uk.RSS genus *Brachyactis* Ledeb. Ukr. Bot. Journ. **43**, 2. Pp. 51–54].

ПРОТОПОПОВА В.В. Lycopodiophyta, Equisetophyta, Polypodiophyta, *Orchidaceae* // Хорология флоры Украины / Отв. ред. Е.Н. Кондратюк. – Киев: Наук. думка, 1986. – С. 17–39, 156–163, 166–169, 172, 175, 245–250 [PROTOROVA V.V. 1986. Lycopodiophyta, Equisetophyta, Polypodiophyta, *Orchidaceae*. Chorology of Ukrainian flora. Ed. E.N. Kondratyuk. Pp. 17–39, 156–163, 166–169, 172, 175, 245–250. Kyiv: Naukova dumka Press].

ПРОТОПОПОВА В.В. Систематична структура та флорогенетичні зв'язки синантропної флори України // Укр. ботан. журн. – 1986. – **43**, № 3. – С. 40–45 [PROTOROVA V.V. 1986. Systematic structure and phylogenetic connection of synanthropic flora of Ukraine. Ukr. Bot. Journ. **43**, 3. Pp. 40–45].

МОРОЗЮК С.С., ПРОТОПОПОВА В.В. Трав'янисті рослини. 2 вид., доповн. – К.: Рад. шк., 1986. – 159 с. [MOROZYUK S.S., PROTOROVA V.V. 1986. Herbaceous plants. 2 ed. 159 p. Kyiv: Radyanska shkola Press.]

#### 1987

ПРОТОПОПОВА В.В. Флористичні комплекси синантропної флори України // Укр. ботан. журн. – 1987. – **44**, № 3.– С. 36–41 [PROTOROVA V.V. 1987. Floristic complexes of synanthropic flora of Ukraine. Ukr. Bot. Journ. **44**, 3. Pp. 36–41].

ПРОТОПОПОВА В.В. Lycopodiophyta, Equisetophyta, Polypodiophyta, *Nyctaginaceae*, *Amaranthaceae* *Oxalidaceae*, *Balsaminaceae*, *Martiniaceae*, *Asteraceae* (*Ambrosia* L., *Bidens* L., *Cyclachaena* Fresen., *Galinsoga* Ruiz. et Pav., *Grindelia* Willd., *Xanthium* L., *Ximenesia* Cav.), *Amarilidaceae*, *Dioscoriaceae*, *Pontederiaceae*, *Iridaceae*, *Orchidaceae* // Определитель высших растений / Отв. ред. Ю.Н. Прокудин. – Киев: Наук. думка, 1987. – С. 21–37, 64, 83–84, 219–220, 223, 294, 320, 328–331, 401–412, 470 [PROTOROVA V.V. 1987. Lycopodiophyta, Equisetophyta, Polypodiophyta, *Nyctaginaceae*, *Amaranthaceae* *Oxalidaceae*, *Balsaminaceae*, *Martiniaceae*, *Asteraceae* (*Ambrosia* L., *Bidens* L., *Cyclachaena* Fresen., *Galinsoga* Ruiz. et Pav., *Grindelia* Willd., *Xanthium* L., *Ximenesia* Cav.), *Amarilidaceae*, *Dioscoriaceae*, *Pontederiaceae*, *Iridaceae*, *Orchidaceae* // Determinator of vascular plants of Ukraine. Ed. Yu.N. PROKUDIN. Pp. 21–37, 64, 83–84, 219–220, 223, 294, 320, 328–331, 401–412, 470. Kyiv: Naukova dumka Press].

ПАНОВА Л.С., ПРОТОПОПОВА В.В. Весняні рослини. – К.: Рад. шк., 1987. – 159 с. [PANOVA L.S., PROTOROVA V.V. 1987. Spring plants. 159 p. Kyiv: Radyanska shkola Press].

#### 1988

ПРОТОПОПОВА В.В. Натуралізація адвентивних рослин України // Укр. ботан. журн. – 1988. – **45**, № 4.– С. 10–15 [PROTOROVA V.V. 1988. Naturalization of alien plants in Ukraine. Ukr. Bot. Journ. **45**, 4. Pp. 10-15].

#### 1989

ПРОТОПОПОВА В.В. Особенности натурализации адвентивных растений на Украине // Мат-лы совещ. „Проблемы изучения адвентивной флоры СССР (г. Москва, 1-3 февраля 1989 г.). – М.: Наука. 1989. – С. 59-60 [PROTOROVA V.V. 1989. Peculiarities of naturalization of alien plants in Ukraine. Math. of Conf. „Problem of study of alien plants of USSR” (Moscow, 1-3 february 1989). Pp. 59-60. Moscow: Nauka Press].

ПРОТОПОПОВА В.В. Рослини-мандрівники. – К.: Рад. шк., 1989. – 238 с. [PROTOROVA V.V. 1989. Plants-travellers. 238 p. Kyiv: Radyanska shkola Press].

#### 1991

ПРОТОПОПОВА В.В. Синантропная флора Украины и пути ее развития. – Киев: Наук. думка, 1991. – 204 с. [PROTOROVA V.V. Synanthropic flora of Ukraine and ways of its development. Ed. D. DOBROCHAYEVA. 204 p. Kyiv: Naukova dumka Press].

#### 1994

ПРОТОПОПОВА В.В. Семейство *Asteraceae* (триба *Heliantheae* Cass., кроме рода *Arnica* L.) // Флора европейской части СССР / Под. ред. Н.Н. ЦВЕЛЕВА. – Л.: Наука, 1994. – Т. 7. – С. 25–52 [PROTOROVA V.V. Family *Asteraceae* (tribe *Heliantheae* Cass., excl. *Arnica* L.). Flora of European part of USSR. Ed. N.N. TZVELEV – Vol. 7. Pp. 25-52. Leningrad, Nauka Press].

ПРОТОПОПОВА В.В., МОСЯКІН С.Л., ШЕВЕРА М.В. *Lepidium virginicum* L. – новий вид флори України // Укр. ботан. журн. – 1994. – 51, № 2/3. – С. 225–228 [PROTOROVA V.V., MOSYAKIN S.L., SHEVERA M.V. 1996. *Lepidium virginicum* L. – new species of Ukrainian flora. Ukr. Bot. Journ. 51, 2-3. Pp. 225-228].

#### 1996

ПРОТОПОПОВА В.В. Родина *Orchidaceae* // Червона книга України. Рослинний світ. 2 вид. / Під заг. ред. Ю.Р. ШЕЛЯГА-СОСОНКА. – К.: УЕ, 1996. – С. 338–354, 460–403 [PROTOROVA V.V. 1996. Red data Book of Ukraine. Ed. YU.R. SHELYAG-SOSONKO. Pp. 338–354, 460–403. Kiev: Ukrainian encyclopedia].

PROTOROVA V. 1996. Trend of synanthropization of the flora of rural settlements and their buffer zones in the lower reaches of the Southern Bug river // Proceeding of II Intern. Conf. Anthropization and environment of rural settlements. Flora and vegetation (Tarczal-Tokaj, 24-28 July 1996) / Ed. by A. TERPO and C. MOCHNASKY. – Pp. 55–58.

#### 1998

ПРОТОПОПОВА В.В. Адвентизація флори України (проблеми та перспективи вирішення) // Мат-ли Міжнар. конф. Промислова ботаніка (м. Донецьк, 1998 р.). – Донецьк, 1998. – С. 76–81 [PROTOROVA V.V. Adventization of Ukrainian flora (problems and perspectives of decision). Math. of Intern. Conf. Industrial botany (Donetsk, 1998). Pp. 76–81].

PROTOROVA V. Ecotopological differentiation of Kyiv ruderal flora // II Inter. Conf. Anthropization and environmental of rural settlements. Flora and vegetation (Zemplinska Sirava, 1998). – Zemplinska Sirava, 1998. – Pp. 49–52.

PROTOROVA V., SHEVERA M.V. 1998. Expansion of alien plants in riverside in Tisa river basin (Transcarpathia) // Thaiszia. – J. Bot. Kosice. – Vol. 8, 2. – Pp. 33–42.

#### 1999

PROTOROVA V., SHEVERA M.V. 1999. Analysis of the modern phytoinvasions in Ukraine // Proceedings 5 Intern. Conf. on the ecology of invasive alien plants (Maddalena, 1999). – Pp. 94–95.

#### 2000

Дідух Я.П., Плюта П.Г., ПРОТОПОПОВА В.В., Єрмоленко В.М., Коротченко І.А., Каркуцієв Г.М., Бурда Р.І. Екофлора України. – Т. 1 / Відпов. ред. Я.П. Дідух. – К.: Фітосоціоцентр, 2000. – 284 с. [DIDUKH YA.P., PLYUTA G.G., PROTOROVA V.V., YERMOLENKO V.M., KOROTCHENKO I.A., KARKUTSIYEV G.M., BURDA R.I. 2000. Ecoflora of Ukraine. Ed. Didukh Ya.P. – Vol. 1. 284 p. – Kiev: Phytosociocenter Press].

Крицька Л.І., ПРОТОПОПОВА В.В., ФЕДОРОНЧУК М.М., ШЕВЕРА М.В. Типіфікація видів судинних рослин, описаних з території України: родини *Solanaceae* Juss., *Scrophulariaceae* Juss., *Orobanchaceae* Vent., *Lentibulariaceae* Rich., *Plantaginaceae* Juss. // Укр. ботан. журн. – 2000. – 57, 5. – С. 533–541 [KRYTZKA L.I., PROTOROVA V.V., FEDORONCHUK M.M., SHEVERA M.V. 2000. Typification of species of vascular plants described from Ukraine: families *Solanaceae* Juss.,

*Scrophulariaceae* Juss., *Orobanchaceae* Vent., *Lentibulariaceae* Rich., *Plantaginaceae* Juss. Ukr. Bot. Journ. **57**, 5. Pp. 533–541].

#### 2002

- ПРОТОПОПОВА В.В., ГРИГОРАК М.Ю., ШЕВЕРА М.В. Еколого-економічні і логістичні аспекти фітозабруднення в Україні // Регіонал. перспективи. – 2002. – 2 (21). – С. 19–21 [PROTOROVA V.V., GRIGORAK M.J., SHEVERA M.V. 2002. Ecological, economic and logistic aspects of biopollution in Ukraine. Regional perspectives. **2** (21). Pp. 19–21].
- ПРОТОПОПОВА В.В., МОСЯКІН С.Л., ШЕВЕРА М.В. Фітоінвазії в Україні як загроза біорізноманіттю: сучасний стан і завдання на майбутнє. – К.: Ін-тут ботаніки НАНУ, 2002. – 32 с. [PROTOROVA V.V., MOSYAKIN S.L., SHEVERA M.V. 2002. Plant invasions in Ukraine as a threat to biodiversity: the present situation and tasks for the future. Kiev. Institute of botany, NAS of Ukraine. 32 p.].
- PROTOROVA V.V. 2002. The Mediterranean element in the alien flora of Ukraine // V Intern. Conf. Anthropization and environment of rural settlements. Flora and vegetation (Uzhgorod, Kostryno, 16-18 May 2002). – Kyiv: Phytosociocenter Press. – Abstracts. – Pp. 71–72.
- PROTOROVA V., SHEVERA M. 2002. A preliminary checklist of urban flora of Uzhgorod. – 68 p. Kyiv: Phytosociocenter Press.
- SHEVERA M., KUZMICHENOV A., PROTOROVA V., MOSYAKIN S. 2002. Comparative urban floristics in Eastern and Central Europe: time to joint forces? // V Intern. Conf. Anthropization and environment of rural settlements. Flora and vegetation (Uzhgorod, Kostryno, 16-18 May 2002). – Pp. 80-81. Kyiv: Phytosociocenter Press.

#### 2003

- ПРОТОПОПОВА В.В., МОСЯКІН С.Л., ШЕВЕРА М.В. Вплив адвентивних видів рослин на фітобіоту України // Оцінка і напрямки зменшення загроз біорізноманіття / Відпов. ред. О. Дудкін. – К.: Хімджест, 2003. – С. 129–155, 358, 364–374 [PROTOROVA V.V., MOSYAKIN S.L., SHEVERA M.V. 2003. Influence of alien plants on phytobiota of Ukraine. Estimation and directions of decreasing of threats of biodiversity. Ed. O. DUDKIN. – Pp. 129–155, 358, 364–374. Kyiv: Khimdzhest Press].
- PROTOROVA V. 2003. Phytogeographical characteristics of stable and non-stable alien components of the Kyiv urban flora // Phytogeographical problems of synanthropic plants / Ed. A. ZAJAC, M. ZAJAC, B. ZEMANEK. – Cracow, Institute of Botany Jagiellonian university. – P. 41-46.

#### 2004

- ГРИГОРАК М.Ю., ПРОТОПОПОВА В.В., ШЕВЕРА М.В. Аспекти економічної оцінки впливу неаборигенних рослин України на довкілля // Регіонал. перспективи. – 2004. – № 6 (43). – С. 44-48. [GRIGORAK M.J., PROTOROVA V.V., SHEVERA M.V. Aspects of economic estimation of influence of alien plants of Ukraine on environment. Regional perspectives. **6** (43). Pp. 44–48].
- КАГАЛО О.О., ПРОТОПОПОВА В.В., ШЕВЕРА М.В. Біорізноманіття в урбанізованому ландшафті: особливості формування і проблеми оцінки // Біорізноманіття Кам'янця-Подільського. Попередній критичний інвентаризаційний конспект рослин, грибів і тварин / За ред. О.О. КАГАЛО, М.В. ШЕВЕРИ, А.А. ЛЕВАНЦЯ. – Львів: Ліга-Прес, 2004. – С. 10–16. [KAGALO O.O., PROTOROVA V.V., SHEVERA M.V. Biodiversity in urban landscape: peculiarities of forming and problems of estimation In Biodiversity of Kamyanets-Podilsky. A preliminary critical inventarization checklist

of plants, fungi, and animals. Ed. KAGALO O.O., SHEVERA M.V., LEVANETS A.A. Pp. 10-16. Lviv: Liga-Press].

КАГАЛО О.О., СКІБІЦЬКА Н.В., ЛЮБІНСЬКА Л.Г., ГУЗІК Я., ПРОТОПОПОВА В.В., ШЕВЕРА М.В. Судинні рослини м. Кам'янець-Подільський // Біорізноманіття Кам'янця-Подільського. Попередній критичний інвентаризаційний конспект рослин, грибів і тварин / За ред. О.О. КАГАЛО, М.В. ШЕВЕРИ, А.А. ЛЕВАНЦЯ. – Львів: Ліга-Прес, 2004. – С. 82–134 [KAGALO O.O., SKIBYNSKA N.V., LYUBINSKA, GUZIK J., PROTODOROVA V.V., SHEVERA M.V. 2004. Vascular plants of Kamyanets-Podilsky In Biodiversity of Kamyanets-Podilsky. A preliminary critical inventarization checklist of plants, fungi, and animals. Ed. KAGALO O.O., SHEVERA M.V., LEVANETS A.A. Pp. 82-134. Lviv: Liga-Press].

ФЕДОРОНЧУК М.М., АНДРІЄНКО Т.Л., ПРОТОПОПОВА В.В. Про необхідність створення НПП Переяславський, як складової екологічної мережі України // Молодь у вирішенні регіональних та транскордонних проблем екологічної безпеки. Перспективи формування Пан'європейської екологічної мережі: Мат-ли третьої міжнар. наук. конф. (м. Чернівці, 21-22 квітня 2004 р.). – Чернівці: Зелена Буковина, 2004. – С. 99–105. [FEDORONCHUK N.M., ANDRIYENKO T.L., PROTODOROVA V.V. 2004. About necessity of creation of NNP Pereyaslavsky as component of ecological network of Ukraine. Youth in decision of regional and trans-boundary problems of ecological security. Perspectives of forming Pan-European econet: Math. of III intern. conf. (Chernivtsi, 21-22 April 2004). Chernivtsi: Zelena Bucovyna Press. Pp.99-105].

### 2005

ПРОТОПОПОВА В.В., ШЕВЕРА М.В. Фітоінвазії. I. Аналіз основних термінів // Промышленная ботаника. – 2005. – Вип. 5. – С. 55–60 [PROTODOROVA V., SHEVERA M. 2005. Phytoinvasions. I. Analysis of main terms. Industrial Botany. Vol. 5. Pp. 55-60].

ПРОТОПОПОВА В. 2005. The Mediterranean element of the alien flora of Ukraine // V Intern. Conf. Anthropization and environment of rural settlements. Flora and vegetation. Proceedings / Ed. S.L. MOSYAKIN, M.V. SHEVERA. – Kyiv: M.G. Kholodny Institute of Botany, NAS of Ukraine. – P. 190–203.

ПРОТОПОПОВА В., ШЕВЕРА М. 2005. *Heracleum mantegazzianum* Sommier et Levier in Ukraine // The ecology and management of the Giant Alien *Heracleum mantegazzianum*. Final Intern. Workshop of the „Giant Alien” Project. Programme and Proceedings (Giessen, 21-23.02.2005). – Giessen, 2005. – P. 26.

ПРОТОПОПОВА В., ШЕВЕРА М. 2005. Archaeophytes in Ukraine: the present patterns of distribution and degree of naturalization. Thaiszia. J. Bot. Kosice. – Vol. 15, Suppl. 1. – Pp. 53–69.

### 2006

ПРОТОПОПОВА В.В. Дослідження синантропної флори України: підсумки та перспективи // Синантропізація рослинного покриву України: Тез. наук. допов. (м. Переяслав-Хмельницький, 27-28 квітня 2006 р.). – Переяслав-Хмельницький, 2006. – С. 170–174. [PROTODOROVA V.V. Investigation of synanthropic flora of Ukraine: results and perspectives. Synanthropisation of plant cover of Ukraine. Abstracts. (Pereyaslav-Khmelnitsky, 27-28 april 2006). Pp. 170-174].

ПРОТОПОПОВА В.В., ШЕВЕРА М.В. Адвентивізація природних та штучних екосистем Причорномор'я // Наук. Вісн. НАУ. – 2006. – № 93. – С. 78–88 [PROTODOROVA V.V.,

- SHEVERA M.V. 2006. Adventization of natural and anthropogenic ecosystems of Black Sea area. Visnyk of National Agrarian University. Pp. 78-88].
- ПРОТОРОВА V.V., SHEVERA M.V., MELNYK R.P. 2006. The history of introduction and present distribution of *Elaeagnus angustifolia* L. in the Black Sea area (Ukraine) // Chornomorsky Bot. Journ. – 2, № 2. – С. 1–13.
- ПРОТОРОВА V.V., SHEVERA M.V., MOSYAKIN S.L. 2006. Deliberate and unintentional introduction of invasive weeds: a case study of the alien flora of Ukraine // Euphytica. — 148. – P. 17–33.

#### 2007

- ДЖУРАН В.М., КРЕЦУЛ Н.І., ПРОТОПОЛОВА В.В., ФЕДОРОНЧУК М.М., ШЕВЕРА М.В. Фітозабруднення рослинного покриву Середнього Придніпров'я. Анований конспект синантропної флори. – К.: Автореферат, 2007. – 48 с. [DZHURAN V.M., KRETSUL N.I., PROTODOROVA V.V., FEDORONCHUK N.M., SHEVERA M.V. 2007. Phytomollution of Middle Dnipro plant cover. Annotation checklist of synanthropic flora. 48 p. Kyiv: Avtoreferat Press].
- МОРОЗЮК С.С., ПРОТОПОЛОВА В.В. Трав'янисті рослини України. Атлас-визначник. – Тернопіль: Навчальна книга – Богдан, 2007. – 216 с. [MOROZYUK S.S., PROTODOROVA V.V. 2007. Herbaceous plants of Ukraine. Atlas-manual. 216 p. Ternopil: Navchalna knyga-Bogdan Press].
- ПАНОВА Л.С., ПРОТОПОЛОВА В.В., МОРОЗЮК С.С. Весняні рослини України. – Тернопіль: Навчальна книга – Богдан, 2007. – 160 с. [PANOVA L.S., PROTODOROVA V.V., MOROZYUK S.S. 2007. Spring plants of Ukraine. 160 p. Ternopil: Navchalna knyga-Bogdan Press].

#### 2008

- ПРОТОРОВА V., SHEVERA M. 2008. Participation of archaeophytes in urban floras in different botanical and geographical zones of Ukraine: a preliminary assessment. Thaiszia – J. Bot. Kosice. – Vol. 18, Suppl. 1. – Pp. 89-104.
- ПРОТОРОВА V., SHEVERA M. 2008. Participation of kenophytes in urban floras in different botanical and geographical zones of Ukraine: a preliminary assessment. VIII Internat. Conf. „Anthropization of rural settlements. Flora and vegetation in a changing lanscape”). – Katowice, Poland, 30.06-02.07. 2008). – Katowice, 2008.- Abstracts. – P. 49.
- ПРОТОРОВА V., SHEVERA M. 2008. On the necessity of special study of flora synanthropisation in Econet elements. Neobiota: towards a synthesis.– 5th European Conference on Biological Invasions (Prague, Czech Republic, 23.-26. September 2008). – Ed. P. PYŠEK, J. PERGL. – Prague. – P. 241.