




P. J. ŠAFÁRIK UNIVERSITY
FACULTY OF SCIENCE
INSTITUTE OF MATHEMATICS
Jesenná 5, 040 01 Košice, Slovakia



I. Žežula and D. Klein

**Orthogonal decompositions in growth
curve models**

IM Preprint, series A, No. 4/2010
March 2010

Orthogonal decompositions in growth curve models*

Ivan ŽEŽULA and Daniel KLEIN

Abstract. The article shows the advantage of orthogonal decompositions in the standard and extended growth curve models. Using this, distribution of estimators of ρ and σ^2 in the standard GCM with uniform correlation structure is derived. Also, equivalence of Hu and von Rosen conditions in the extended GCM under mild conditions is shown.

1 The standard GCM with uniform correlation structure

The basic model we consider is the following:

$$Y = XBZ' + e, \quad \text{vec } e \sim N(0, \Sigma \otimes I_n), \quad \Sigma = \theta_1 G + \theta_2 w w', \quad (1)$$

where $Y_{n \times p}$ is a matrix of independent p -variate observations, $X_{n \times m}$ is an ANOVA design matrix, $Z_{p \times r}$ is a regression variables matrix, and e matrix of random errors. As for the unknown parameters, $B_{m \times r}$ is an location parameters matrix, and θ_1, θ_2 are (scalar) variance parameters. Matrix $G_{p \times p} > 0$ and vector $w \in \mathbb{R}^p$ are known. The vec operator stacks elements of a matrix into a vector column-wise. This correlation structure is called generalized uniform correlation structure. It was studied in the context of the growth curve model (GCM) in [6], and recently in [4]. A special case was studied also in [3].

While Žežula in [6] used directly model (1), Ye and Wang used modified model with orthogonal decomposition:

$$\begin{aligned} Y &= Y_1 + Y_2, \\ Y_1 &= YG^{-\frac{1}{2}}P_F = XBZ'G^{-\frac{1}{2}}P_F + e_1, \\ Y_2 &= YG^{-\frac{1}{2}}M_F = XBZ'G^{-\frac{1}{2}}M_F + e_2, \end{aligned} \quad (2)$$

*The research was supported by grants VEGA MŠ SR 1/0131/09 and VEGA MŠ SR 1/0325/10.

where $F = G^{-\frac{1}{2}}w$, P_F is the orthogonal projection matrix onto the column space $\mathcal{R}(F)$ of F , and $M_F = I - P_F$ onto its orthogonal complement.

Let us denote

$$S = \frac{1}{n - r(X)} Y' M_X Y,$$

$$W_1 = P_F G^{-\frac{1}{2}} S G^{-\frac{1}{2}} P_F, \quad W_2 = M_F G^{-\frac{1}{2}} S G^{-\frac{1}{2}} M_F.$$

The estimators of Žežula are

$$\hat{\theta}_1 = \frac{(\mathbf{1}'w)^2 \text{Tr}(S) - \mathbf{1}'S\mathbf{1} w'w}{(\mathbf{1}'w)^2 \text{Tr}(G) - \mathbf{1}'G\mathbf{1} w'w}, \quad \hat{\theta}_2 = \frac{\mathbf{1}'S\mathbf{1} \text{Tr}(G) - \mathbf{1}'G\mathbf{1} \text{Tr}(S)}{(\mathbf{1}'w)^2 \text{Tr}(G) - \mathbf{1}'G\mathbf{1} w'w}, \quad (3)$$

and the estimators of Ye and Wang are

$$\hat{\theta}_1^* = \frac{\text{Tr}(W_2)}{p-1} = \frac{w'G^{-1}w \cdot \text{Tr}(G^{-1}S) - w'G^{-1}SG^{-1}w}{(p-1)(w'G^{-1}w)},$$

$$\hat{\theta}_2^* = \frac{(p-1)\text{Tr}(W_1) - \text{Tr}(W_2)}{(p-1)w'G^{-1}w} = \frac{p \cdot w'G^{-1}SG^{-1}w - w'G^{-1}w \cdot \text{Tr}(G^{-1}S)}{(p-1)(w'G^{-1}w)^2}. \quad (4)$$

These pairs of estimators are both unbiased, but different. Naturally, we would like to know the variances. Since $S \sim \mathcal{W}_p\left(n - r(X), \frac{1}{n - r(X)}\Sigma\right)$, it is easy to establish that

$$\text{var}(\text{vec } S) = \frac{1}{n - r(X)} (I_{p^2} + K_{pp}) (\Sigma \otimes \Sigma), \quad (5)$$

where K_{pp} is the commutation matrix, see e.g. [5]. This immediately implies

$$\text{var } \text{Tr}(G^{-1}S) = \frac{2}{n - r(X)} \text{Tr}(G^{-1}\Sigma G^{-1}\Sigma),$$

$$\text{var}(w'G^{-1}SG^{-1}w) = \frac{2}{n - r(X)} (w'G^{-1}\Sigma G^{-1}w)^2,$$

$$\text{cov}[\text{Tr}(G^{-1}S), w'G^{-1}SG^{-1}w] = \frac{2}{n - r(X)} w'G^{-1}\Sigma G^{-1}\Sigma G^{-1}w.$$

Necessary formulas for $\text{var } \text{Tr}(S)$, $\text{var } \mathbf{1}'S\mathbf{1}$, and $\text{cov}(\text{Tr}(S), \mathbf{1}'S\mathbf{1})$ are special cases. Short computation gives

$$\text{var } \hat{\theta}_1 = \frac{2}{n - r(X)} \cdot \frac{(\mathbf{1}'w)^4 \text{Tr}(\Sigma^2) - 2(\mathbf{1}'w)^2 w'w \cdot \mathbf{1}'\Sigma^2\mathbf{1} + (w'w)^2 (\mathbf{1}'\Sigma\mathbf{1})^2}{[(\mathbf{1}'w)^2 \text{Tr}(G) - \mathbf{1}'G\mathbf{1} w'w]^2}, \quad (6)$$

$$\text{var } \hat{\theta}_2 = \frac{2}{n - r(X)} \cdot \frac{[\text{Tr}(G)\mathbf{1}'\Sigma\mathbf{1}]^2 - 2\text{Tr}(G)\mathbf{1}'G\mathbf{1}\mathbf{1}'\Sigma^2\mathbf{1} + (\mathbf{1}'G\mathbf{1})^2 \text{Tr}(\Sigma^2)}{[(\mathbf{1}'w)^2 \text{Tr}(G) - \mathbf{1}'G\mathbf{1} w'w]^2}, \quad (7)$$

and

$$\text{var } \hat{\theta}_1^* = \frac{2}{n-r(X)} \cdot \frac{1}{(p-1)^2 (w'G^{-1}w)^2} \left[(w'G^{-1}w)^2 \text{Tr}(G^{-1}\Sigma G^{-1}\Sigma) + \right. \\ \left. + (w'G^{-1}\Sigma G^{-1}w)^2 - 2(w'G^{-1}w)(w'G^{-1}\Sigma G^{-1}\Sigma G^{-1}w) \right], \quad (8)$$

$$\text{var } \hat{\theta}_2^* = \frac{2}{n-r(X)} \cdot \frac{1}{(p-1)^2 (w'G^{-1}w)^4} \left[(w'G^{-1}w)^2 \text{Tr}(G^{-1}\Sigma G^{-1}\Sigma) + \right. \\ \left. + p^2 (w'G^{-1}\Sigma G^{-1}w)^2 - 2p(w'G^{-1}w)(w'G^{-1}\Sigma G^{-1}\Sigma G^{-1}w) \right]. \quad (9)$$

Analytical comparison of these quantities is quite difficult. Few simulations performed suggest that in general Ye and Wang's estimators tend to have smaller variance.

Very important special case of the previous model is the model with

$$\Sigma = \sigma^2 [(1-\rho)I_p + \rho \mathbf{1}\mathbf{1}']. \quad (10)$$

This correlation structure is called the uniform correlation structure or the intraclass correlation structure. It is the case with $G = I_p$ and $w = \mathbf{1}$, slightly reparametrized. It must hold

$$-\frac{1}{p-1} \leq \rho \leq 1.$$

As a special case of (1), estimators of σ^2 and ρ can be then obtained by a simple transformation of $\hat{\theta}_1$ and $\hat{\theta}_2$:

$$\hat{\sigma}^2 = \hat{\theta}_1 + \hat{\theta}_2 \quad \text{and} \quad \hat{\rho} = \frac{\hat{\theta}_2}{\hat{\theta}_1 + \hat{\theta}_2}. \quad (11)$$

This implies the following form of estimators due to Žežula:

$$\hat{\sigma}_Z^2 = \frac{\text{Tr}(S)}{p}, \quad \hat{\rho}_Z = \frac{1}{p-1} \left(\frac{\mathbf{1}'S\mathbf{1}}{\text{Tr}(S)} - 1 \right), \quad (12)$$

and due to Ye and Wang:

$$\hat{\sigma}_{YW}^2 = \frac{\text{Tr}(V_1) + \text{Tr}(V_2)}{p}, \quad \hat{\rho}_{YW} = 1 - \frac{p \text{Tr}(V_2)}{(p-1)(\text{Tr}(V_1) + \text{Tr}(V_2))}, \quad (13)$$

where

$$V_1 = P_1 S P_1, \quad V_2 = M_1 S M_1.$$

Ye and Wang recognized that $\hat{\sigma}_Z^2 = \hat{\sigma}_{YW}^2$, but they failed to recognize that also $\hat{\rho}_Z = \hat{\rho}_{YW}$.

Lemma 1. $\hat{\sigma}_Z^2 = \hat{\sigma}_{YW}^2$ and $\hat{\rho}_Z = \hat{\rho}_{YW}$ for any Y .

Proof. Trivially,

$$\text{Tr}(V_1) = \text{Tr}(P_1 S P_1) = \text{Tr}(S P_1) = \frac{1}{p} \text{Tr}(S \mathbf{1} \mathbf{1}') = \frac{1}{p} \mathbf{1}' S \mathbf{1},$$

and

$$\text{Tr}(V_2) = \text{Tr}(M_1 S M_1) = \text{Tr}(S M_1) = \text{Tr}(S) - \text{Tr}(S P_1) = \text{Tr}(S) - \frac{1}{p} \mathbf{1}' S \mathbf{1}.$$

Substituting these values into (13), we easily get (12). \square

Thus, in the following we can write only $\hat{\sigma}^2$ and $\hat{\rho}$.

This orthogonal decomposition is very useful for derivation of the distribution of the estimators.

Lemma 2. *Let $H \sim \mathcal{W}_p(\ell, \Xi)$, $\Xi > 0$, and $T_{k \times p}$ be arbitrary matrix. Then,*

$$\text{Tr}(THT') \sim \sum_{i=1}^{r(T)} \lambda_i \chi_{\ell}^2,$$

where $\lambda_1, \dots, \lambda_{r(T)}$ are all positive eigenvalues of $T\Xi T'$ and $r(T)$ is the rank of T . In particular,

$$\text{E Tr}(THT') = \ell \sum_{i=1}^{r(T)} \lambda_i, \quad \text{var Tr}(THT') = 2\ell \sum_{i=1}^{r(T)} \lambda_i^2.$$

Proof. There must exist independent r.v. X_1, \dots, X_{ℓ} distributed as $N_p(0, \Xi)$ such that $H = \sum_{i=1}^{\ell} X_i X_i' = G'G$, where $G' = (X_1, \dots, X_{\ell})$. Then, $TX_i \sim N_k(0, T\Xi T') \forall i$ (which may be singular). According to the Theorem in [1] it holds

$$\text{Tr}(THT') = \text{Tr}((GT')'(GT')) \sim \sum_{i=1}^{r(T)} \lambda_i \chi_{\ell}^2,$$

where $\lambda_1, \dots, \lambda_{r(T)}$ are all positive eigenvalues of $T\Xi T'$. Since χ^2 's are independent, the claims about mean and variance are trivial. \square

The following results concerning distributions of $\text{Tr}(V_1)$ and $\text{Tr}(V_2)$ can be found in [4], but without proof. We extend these results to the parameters of interest.

Theorem 3. *Distributions of $\text{Tr}(V_1)$ and $\text{Tr}(V_2)$ are independent,*

$$\begin{aligned} \text{Tr}(V_1) &\sim \frac{\sigma^2[1 + (p-1)\rho]}{n - r(X)} \chi_{n-r(X)}^2, \\ \text{Tr}(V_2) &\sim \frac{\sigma^2(1 - \rho)}{n - r(X)} \chi_{(p-1)(n-r(X))}^2, \end{aligned}$$

so that

$$\hat{\sigma}^2 \sim \frac{\sigma^2}{p(n-r(X))} \left[(1 + (p-1)\rho)\chi_{n-r(X)}^2 + (1-\rho)\chi_{(p-1)(n-r(X))}^2 \right],$$

$$\frac{1-\rho}{1+(p-1)\rho} \left[\frac{1+(p-1)\hat{\rho}}{1-\hat{\rho}} \right] \sim F_{n-r(X), (p-1)(n-r(X))}.$$

Proof. It is well known that under normality

$$S \sim \mathcal{W}_p \left(n-r(X), \frac{1}{n-r(X)} \Sigma \right).$$

(see e.g. Theorem 3.8 in [5]). We want to make use of Lemma 2 with $\ell = n-r(X)$, $\Xi = \frac{1}{n-r(X)} \Sigma$, $T = P_1$, and also with $T = M_1$. Since

$$P_1 \left(\frac{1}{n-r(X)} \Sigma \right) P_1 = \frac{\sigma^2[1+(p-1)\rho]}{n-r(X)} P_1$$

is a multiple of idempotent matrix of rank 1, its only positive eigenvalue is equal to $\sigma^2[1+(p-1)\rho]/(n-r(X))$. Similarly, since M_1 is idempotent with rank $p-1$ and

$$M_1 \left(\frac{1}{n-r(X)} \Sigma \right) M_1 = \frac{\sigma^2(1-\rho)}{n-r(X)} M_1,$$

it has $p-1$ positive eigenvalues which are all equal to $\sigma^2(1-\rho)/(n-r(X))$. Now the results for $\text{Tr}(V_1)$ and $\text{Tr}(V_2)$ follow from Lemma 2, perpendicularity of M_1 and P_1 , and properties of χ^2 -distribution.

This, together with (13), immediately implies the result for $\hat{\sigma}^2$. The second formula in (13) can be transformed to

$$\frac{1+(p-1)\hat{\rho}}{1-\hat{\rho}} = \frac{(p-1) \text{Tr}(V_1)}{\text{Tr}(V_2)}.$$

Because the distributions of $\text{Tr}(V_1)$ and $\text{Tr}(V_2)$ are independent, clearly

$$\frac{\sigma^2(1-\rho)(n-r(X))}{\sigma^2[1+(p-1)\rho](n-r(X))} \cdot \frac{(p-1) \text{Tr}(V_1)}{\text{Tr}(V_2)} \sim F_{n-r(X), (p-1)(n-r(X))}.$$

□

This result is not very useful with respect to $\hat{\sigma}^2$, since its distribution depends on both σ^2 and ρ , but enables us to test for any specific value of ρ . Using simple

transformation, we can even derive directly probability density function of $\hat{\rho}$:

$$\begin{aligned} f(x) &= \left(\frac{1 - \rho}{(p-1)[1 + (p-1)\rho]} \right)^{\frac{n-r(X)}{2}} \frac{\Gamma\left(\frac{p(n-r(X))}{2}\right)}{\Gamma\left(\frac{n-r(X)}{2}\right) \Gamma\left(\frac{(p-1)(n-r(X))}{2}\right)} \times \\ &\times \left(1 + \frac{1 - \rho}{(p-1)[1 + (p-1)\rho]} \frac{1 + (p-1)x}{1-x} \right)^{-\frac{p(n-r(X))}{2}} \times \\ &\times \left(\frac{1 + (p-1)x}{1-x} \right)^{\frac{n-r(X)}{2} - 1} \frac{p}{(1-x)^2}. \end{aligned}$$

Also, $1 - \alpha$ confidence interval for ρ is given by

$$\left(\frac{1 - c_1}{1 + (p-1)c_1}; \frac{1 - c_2}{1 + (p-1)c_2} \right), \quad (14)$$

where

$$c_1 = \frac{1 - \hat{\rho}}{1 + (p-1)\hat{\rho}} F_{n-r(X), (p-1)(n-r(X))} \left(1 - \frac{\alpha}{2} \right)$$

and

$$c_2 = \frac{1 - \hat{\rho}}{1 + (p-1)\hat{\rho}} F_{n-r(X), (p-1)(n-r(X))} \left(\frac{\alpha}{2} \right).$$

Figures 1–4 below show histograms and theoretical densities of $\hat{\rho}$ for a special case of the model (quadratic growth in three groups, 2500 simulations) for various true values of unknown parameter.





Example: Let us consider random sample from bivariate normal distribution with the same variances in both dimensions. It can be formally written as GCM with the uniform correlation structure:

$$Y = \begin{pmatrix} Y_{11} & Y_{12} \\ \vdots & \vdots \\ Y_{n1} & Y_{n2} \end{pmatrix} = \mathbf{1}_n (\mu_1, \mu_2) I_2 + e, \quad e \sim N_{n \times 2} \left(0_{n \times 2}, \sigma^2 \begin{pmatrix} 1 & \rho \\ \rho & 1 \end{pmatrix} \otimes I_n \right).$$

Using the above mentioned estimator we get

$$\hat{\rho} = \frac{2s_{12}}{s_1^2 + s_2^2},$$

where s_{12} is sample covariance of the two variables, and s_1^2 and s_2^2 sample variances. This estimator is slightly more effective than standard sample correlation coefficient (in the sense of MSE).

FIGURE 1. $\rho = -0,2$ FIGURE 2. $\rho = 0$ FIGURE 3. $\rho = 0,5$ FIGURE 4. $\rho = 0,95$

2 The extended growth curve model

The extended growth curve model (EGCM) with fixed effects, called also sum-of-profiles model, is

$$Y = \sum_{i=1}^k X_i B_i Z_i' + e, \quad e \sim N_{n \times p}(0, \Sigma \otimes I_n). \quad (15)$$

The dimensions of matrices X_i , B_i , and Z_i are $n \times m_i$, $m_i \times r_i$, $p \times r_i$, respectively. Usually it is supposed that column spaces of X_i 's are ordered,

$$\mathcal{R}(X_k) \subseteq \dots \subseteq \mathcal{R}(X_1), \quad (16)$$

while nothing is said about different Z_i 's. Recently, Hu (see [2]) came up with modification of the model, assuming

$$X_i' X_j = 0 \quad \forall i \neq j. \quad (17)$$

His idea is to separate groups rather than models. We will show that the two models are under certain conditions equivalent.

Example: Let us consider EGCM with two groups with different growth patterns – linear and quadratic:

$$\begin{aligned} Y_{ij} &= \beta_1 + \beta_2 t_j + e_{ij}, & i &= 1, \dots, n_1, \quad j = 1, \dots, p, \\ &= \beta_3 + \beta_4 t_j + \beta_5 t_j^2 + e_{ij}, & i &= n_1 + 1, \dots, n_1 + n_2, \quad j = 1, \dots, p. \end{aligned}$$

This model can be written as

$$Y = \begin{pmatrix} \mathbf{1}_{n_1} & 0 \\ 0 & \mathbf{1}_{n_2} \end{pmatrix} \begin{pmatrix} \beta_1 & \beta_2 \\ \beta_3 & \beta_4 \end{pmatrix} \begin{pmatrix} 1 & \dots & 1 \\ t_1 & \dots & t_p \end{pmatrix} + \begin{pmatrix} 0 \\ \mathbf{1}_{n_2} \end{pmatrix} \beta_5 (t_1^2 \dots t_p^2) + e,$$

or, by the new way, as

$$Y = \begin{pmatrix} \mathbf{1}_{n_1} \\ 0 \end{pmatrix} (\beta_1, \beta_2) \begin{pmatrix} 1 & \dots & 1 \\ t_1 & \dots & t_p \end{pmatrix} + \begin{pmatrix} 0 \\ \mathbf{1}_{n_2} \end{pmatrix} (\beta_3, \beta_4, \beta_5) \begin{pmatrix} 1 & \dots & 1 \\ t_1 & \dots & t_p \\ t_1^2 & \dots & t_p^2 \end{pmatrix} + e.$$

Note that in the second form in the previous example $\mathcal{R}(Z_1) \subset \mathcal{R}(Z_2)$. This leads to idea that we can consider a model in which the column spaces of all Z_i 's are nested, which naturally arises in situations when different groups use polynomial regression functions of different order.

Let us consider model (15) with condition (16), such that

$$n - p \geq \sum_{i=1}^k r(X_i). \quad (18)$$

Since X_i 's are 0-1 matrices whose columns are indicators of different groups, w.l.o.g. we can assume that all columns of X_1 are mutually perpendicular, and columns of every X_{i+1} are a subset of columns of X_i . Let us define $X_k^* = X_k$ and $X_i^* = X_i \setminus X_{i+1}$, $i = 1, \dots, k-1$, where the symbol $X_i \setminus X_{i+1}$ denotes matrix consisting of those columns of X_i which are not in X_{i+1} . It is easy to see, that $X_i = (X_i^*, \dots, X_k^*)$ and $P_{X_i} - P_{X_{i+1}} = P_{X_{i+1}^\perp \cap X_i} = P_{X_i^*}$.

Then, we can reformulate the model (15) with von Rosen's condition (16) in the following way:

$$\begin{aligned} EY &= \sum_{i=1}^k X_i B_i Z_i' = \sum_{i=1}^k (X_i^*, \dots, X_k^*) \begin{pmatrix} B_{ii}^* \\ \vdots \\ B_{ik}^* \end{pmatrix} Z_i' = \sum_{i=1}^k \sum_{j=i}^k X_j^* B_{ij}^* Z_i' = \\ &= \sum_{j=1}^k \sum_{i=1}^j X_j^* B_{ij}^* Z_i' = \sum_{j=1}^k X_j^* (B_{1j}^*, \dots, B_{jj}^*) \begin{pmatrix} Z_1' \\ \vdots \\ Z_j' \end{pmatrix} \stackrel{\text{df}}{=} \sum_{j=1}^k X_j^* B_j^* Z_j^{*'} \quad (19) \end{aligned}$$

(matrices X_j^* have dimensions $n \times m_j^*$ and B_{ij}^* $m_j^* \times r_i$, where $m_i = \sum_{j=i}^k m_j^*$.) It is now easy to see that model (19) satisfies Hu's condition:

$$X_i^{*'} X_j^* = 0 \quad \forall i \neq j. \quad (20)$$

Moreover, now we have

$$Z_i^* = (Z_1, \dots, Z_i), \quad \forall i = 1, \dots, k,$$

which implies $\mathcal{R}(Z_1^*) \subset \cdots \subset \mathcal{R}(Z_k^*)$.

ECGM with Hu's condition is much easier to handle. If all X_i^* 's and Z_i^* 's are of full rank, then all B_i^* 's are estimable, and unbiased LSE \hat{B}_i^* depend only on X_i^* and Z_i^* :

$$\hat{B}_i^* = (X_i^{*'} X_i^*)^{-1} X_i^{*'} Y \Sigma^{-1} Z_i^* (Z_i^{*'} \Sigma^{-1} Z_i^*)^{-1}, \quad (21)$$

see [2]. Such a closed form was difficult to obtain in the von Rosen model. Even for two components the estimators are rather complicated:

$$\begin{aligned} \hat{B}_1 &= (X_1' X_1)^{-1} X_1' Y \Sigma^{-1} Z_1 (Z_1' \Sigma^{-1} Z_1)^{-1} \\ &\quad - (X_1' X_1)^{-1} X_1' P_{X_2} Y \left(P_{Z_2}^{\Sigma^{-1} M_{Z_1}^{\Sigma^{-1}}} \right)' \Sigma^{-1} Z_1 (Z_1' \Sigma^{-1} Z_1)^{-1}, \\ \hat{B}_2 &= (X_2' X_2)^{-1} X_2' Y \Sigma^{-1} Z_2 \left(Z_2' \Sigma^{-1} M_{Z_1}^{\Sigma^{-1}} Z_2 \right)^{-1}, \end{aligned}$$

see [7]. Each \hat{B}_1 and \hat{B}_2 depends on both Z_1 and Z_2 , and \hat{B}_1 even on X_2 .

Estimator of common variance matrix can be split into perpendicular pieces:

$$\hat{\Sigma} = \frac{1}{n - r(X_1)} Y' M_{X_1} Y = \frac{1}{n - \sum_{i=1}^k r(X_i^*)} Y' \left(I - \sum_{i=1}^k P_{X_i^*} \right) Y. \quad (22)$$

In the last expression, the left-hand side term is the estimator using von Rosen's model and right-hand side one using Hu's model. It is easy to see, that the estimators are equivalent, since $X_1 = (X_1^*, \dots, X_k^*)$.

The situation in Hu's model is much easier also for a special correlation structure $\Sigma = \sigma^2 R$ with R known. The unbiased estimator of residual variance σ^2 is

$$\hat{\sigma}^2 = \frac{1}{n - \sum_{i=1}^k r(X_i) \text{Tr}(P_{Z_i}^{R^{-1}} R)} \text{Tr} \left((Y - \hat{Y})' (Y - \hat{Y}) \right),$$

where $\hat{Y} = \sum_{i=1}^k P_{X_i} Y \left(P_{Z_i}^{\Sigma^{-1}} \right)'$ is the unbiased estimator of EY .

3 Conclusions

Method of orthogonal decomposition is very promising in complex models. Many tasks, which are very difficult or impossible to handle in basic models, can be done with ease in models consisting of mutually orthogonal components. As it is shown above, simple transformation can change a model into an equivalent which allows to determine explicit forms of estimators and/or their distribution. We hope the method will prove even more useful in the future, whether in the models investigated here or in some others.

References

- [1] Glueck, D.H., Muller, K.E. (1998). On the trace of a Wishart. *Communications in Statistics - Theory and Methods* **27(9)**, 2137–2141.
- [2] Hu, J. (2009). Properties of the Explicit Estimators in the Extended Growth Curve Model. *Statistics*, DOI 10.1080/02331880903236884 .
- [3] Klein, D., Žežula, I. (2007). On uniform correlation structure. *Mathematical Methods In Economics And Industry, Herľany, Slovakia*, proceedings, 94–100.
- [4] Ye, R.D., Wang, S.G. (2009). Estimating parameters in extended growth curve models with special covariance structures. *Journal of Statistical Planning and Inference* **139**, 2746–2756.
- [5] Žežula, I. (1993). Covariance components estimation in the growth curve model. *Statistics* **24**, 321–330.
- [6] Žežula, I. (2006). Special variance structures in the growth curve model. *Journal of multivariate analysis* **97**, 606–618.
- [7] Žežula, I. (2008). Remarks on unbiased estimation of the sum-of-profiles model parameters. *Tatra Mountains Mathematical Publications* **39**, 45–52.

E-mail address: ivan.zezula@upjs.sk, daniel.klein@upjs.sk

INSTITUTE OF MATHEMATICS, P. J. ŠAFÁRIK UNIVERSITY, JESENNÁ 5,
SK-04001 KOŠICE, SLOVAKIA

Recent IM Preprints, series A

2006

- 1/2006 Semanišinová I. and Trenkler M.: *Discovering the magic of magic squares*
2/2006 Jendroľ S.: *NOTE – Rainbowness of cubic polyhedral graphs*
3/2006 Horňák M. and Woźniak M.: *On arbitrarily vertex decomposable trees*
4/2006 Cechlárová K. and Lacko V.: *The kidney exchange problem: How hard is it to find a donor ?*
5/2006 Horňák M. and Kocková Z.: *On planar graphs arbitrarily decomposable into closed trails*
6/2006 Biró P. and Cechlárová K.: *Inapproximability of the kidney exchange problem*
7/2006 Rudašová J. and Soták R.: *Vertex-distinguishing proper edge colourings of some regular graphs*
8/2006 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2006*
9/2006 Borbeľová V. and Cechlárová K.: *Pareto optimality in the kidney exchange game*
10/2006 Harminc V. and Molnár P.: *Some experiences with the diversity in word problems*
11/2006 Horňák M. and Zlámalová J.: *Another step towards proving a conjecture by Plummer and Toft*
12/2006 Hančová M.: *Natural estimation of variances in a general finite discrete spectrum linear regression model*

2007

- 1/2007 Haluška J. and Hutník O.: *On product measures in complete bornological locally convex spaces*
2/2007 Cichacz S. and Horňák M.: *Decomposition of bipartite graphs into closed trails*
3/2007 Hajduková J.: *Condorcet winner configurations in the facility location problem*
4/2007 Kovárová I. and Mihalčová J.: *Vplyv riešenia jednej difúznej úlohy a následný rozbor na riešenie druhej difúznej úlohy o 12-tich kockách*
5/2007 Kovárová I. and Mihalčová J.: *Prieskum tvorivosti v žiackych riešeniach vágne formulovanej úlohy*
6/2007 Haluška J. and Hutník O.: *On Dobrakov net submeasures*
7/2007 Jendroľ S., Miškuf J., Soták R. and Škrabuláková E.: *Rainbow faces in edge colored plane graphs*
8/2007 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2007*
9/2007 Cechlárová K.: *On coalitional resource games with shared resources*

2008

- 1/2008 Miškuf J., Škrekovski R. and Tancer M.: *Backbone colorings of graphs with bounded degree*
2/2008 Miškuf J., Škrekovski R. and Tancer M.: *Backbone colorings and generalized Mycielski's graphs*
3/2008 Mojsej I.: *On the existence of nonoscillatory solutions of third order nonlinear differential equations*
4/2008 Cechlárová K. and Fleiner T.: *On the house allocation markets with duplicate houses*

- 5/2008 Hutník O.: *On Toeplitz-type operators related to wavelets*
 6/2008 Cechlárová K.: *On the complexity of the Shapley-Scarf economy with several types of goods*
 7/2008 Zlámalová J.: *A note on cyclic chromatic number*
 8/2008 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2008*
 9/2008 Czap J. and Jendroľ S.: *Colouring vertices of plane graphs under restrictions given by faces*

2009

- 1/2009 Zlámalová J.: *On cyclic chromatic number of plane graphs*
 2/2009 Havet F., Jendroľ S., Soták R. and Škrabuláková E.: *Facial non-repetitive edge-colouring of plane graphs*
 3/2009 Czap J., Jendroľ S., Kardoš F. and Miškuf J.: *Looseness of plane graphs*
 4/2009 Hutník O.: *On vector-valued Dobrakov submeasures*
 5/2009 Haluška J. and Hutník O.: *On domination and bornological product measures*
 6/2009 Kolková M. and Pócsová J.: *Metóda Monte Carlo na hodine matematiky*
 7/2009 Borbeľová V. and Cechlárová K.: *Rotations in the stable b-matching problem*
 8/2009 Mojsej I. and Tartal'ová A.: *On bounded nonoscillatory solutions of third-order nonlinear differential equations*
 9/2009 Jendroľ S. and Škrabuláková E.: *Facial non-repetitive edge-colouring of semiregular polyhedra*
 10/2009 Krajčiová J. and Pócsová J.: *Galtonova doska na hodine matematiky, kvalitatívne určenie veľkosti pravdepodobnosti udalostí*
 11/2009 Fabrici I., Horňák M. and Jendroľ S., ed.: *Workshop Cycles and Colourings 2009*
 12/2009 Hudák D. and Madaras T.: *On local properties of 1-planar graphs with high minimum degree*
 13/2009 Czap J., Jendroľ S. and Kardoš F.: *Facial parity edge colouring*
 14/2009 Czap J., Jendroľ S. and Kardoš F.: *On the strong parity chromatic number*

2010

- 1/2010 Cechlárová K. and Pillárová E.: *A near equitable 2-person cake cutting algorithm*
 2/2010 Cechlárová K. and Jelínková E.: *An efficient implementation of the equilibrium algorithm for housing markets with duplicate houses*
 3/2010 Hutník O. and Hutníková M.: *An alternative description of Gabor spaces and Gabor-Toeplitz operators*